

# Ancient megafaunal mutualisms and extinctions as factors in plant domestication

March 25 2021

---



Muskox (*Ovibos moschatus*) - one of numerous herbivores that roam in the enclosure of the Pleistocene Park, nature reserve in northern Sakha Republic, Russia. This ongoing grazing experiment started in 1988 and its aim is to test if large herbivores can restore rich Pleistocene grassland ecosystems. Credit: Frank Kienast

By clearing forests, burning grasslands, plowing fields and harvesting crops, humans apply strong selective pressures on the plants that survive on the landscapes we use. Plants that evolved traits for long-distance seed-dispersal, including rapid annual growth, a lack of toxins and large seed generations, were more likely to survive on these dynamic anthropogenic landscapes. In a new article, researchers argue that these traits may have evolved as adaptations for megafaunal mutualisms, later allowing those plants to prosper among increasingly sedentary human populations.

The new study hypothesizes that the presence of specific anthropophilic traits explains why a select few [plant families](#) came to dominate the crop and weed assemblages around the globe, such as quinoa, some grasses, and knotweeds. These traits, the authors argue, also explain why so many genera appear to have been domesticated repeatedly in different parts of the world at different times. The 'weediness' and adaptability of those plants was the result of exaptation traits, or changes in the function of an evolutionary trait. In this way, rather than an active and engaged human process, certain plants gradually increased in prominence around villages, in cultivated fields, or on grazing land.

Grasses and field crops weren't the only plants to use prior adaptations to prosper in human landscapes; select handfuls of trees also had advantageous traits, such as large fleshy fruits, resulting from past relationships with large browsers. The rapid extinction of megafauna at the end of the Pleistocene left many of these large-fruited tree species with small, isolated populations, setting the stage for more dramatic changes during later hybridization. When humans began moving these trees they were likely to hybridize with distant relatives, resulting, in some cases, in larger fruits and more robust plants. In this way, the domestication process for many long-generation perennials appears to have been more rapid and tied into population changes due to megafaunal extinctions.



The wood bison (*Bison bison* spp. *athabasca*) of the Ust'-Buotama Bison Park, central Sakha Republic, Russia. In 2006, 35 wood bison were brought from Canada. These megaherbivores adapted to local environment and increased their population. The 100-ha enclosure, where animals live, serves as a study site for ecologists and zoologists and provides an opportunity to trace changes in vegetation associated with herbivore pressure. Credit: Frank Kienast

"The key to better understanding plant domestication may lay further in the past than archaeologists have previously thought; we need to think about the domestication process as another step in the evolution of life on Earth, as opposed to an isolated phenomenon," states Dr. Robert Spengler. He is the director of the archaeobotanical laboratories at the Max Planck Institute for the Science of Human History in Jena,

Germany, and the lead investigator on this paper.

This publication is a result of archaeologists, geneticists, botanists, and paleontologists contributing insights from their unique disciplines to reframe the way scholars think about domestication. The goal of the collaboration is to get researchers to consider the deeper ecological legacies of the [plants](#) and the pre-cultivation adaptations that they study.

Prof. Nicole Boivin, director of the Department of Archaeology at the Max Planck Institute in Jena, studies the ecological impacts of humans deep in the past. "When we think about the ecology of the origins of agriculture, we need to recognize the dramatic changes in plant and animal dynamics that have unfolded across the Holocene, especially those directly resulting from human action," she adds.



While it may just look like a regular field, the plants growing on either side of this bison trail are primarily little barley, one of the North American Lost Crops. This photo was taken during field work by Natalie Muller of Washington University in St. Louis in an attempt to determine the role that bison may have played in shaping the ecology of the progenitors for certain ancient crops. Credit: Natalie Mueller

Ultimately, the scholars suggest that, rather than in archaeological excavations, laboratories, or in modern agricultural fields, the next big discoveries in plant domestication research may come from restored megafaunal landscapes. Ongoing research by Dr. Natalie Mueller, one of the authors, on North American restored prairies is investigating potential links between bison and the North American Lost Crops. Similar studies could be conducted on restored megafaunal landscapes in Europe, such as the Bialowieski National Park in Poland, the Ust'-Buioma Bison Park or Pleistocene Park in Sakha Republic, Russia.

Dr. Ashastina, another author on the paper and a paleontologist studying Pleistocene vegetation communities in North Asia, states, "these restored [nature preserves](#) provide a novel glimpse deep into the nature of plant and animal interactions and allow ecologists, not only to directly trace vegetation changes occurring under herbivore pressure in various ecosystems, but to disentangle the deeper legacies of these mutualisms."

**More information:** Robert N. Spengler et al, Exaptation Traits for Megafaunal Mutualisms as a Factor in Plant Domestication, *Frontiers in Plant Science* (2021). [DOI: 10.3389/fpls.2021.649394](https://doi.org/10.3389/fpls.2021.649394)

Provided by Max Planck Society

Citation: Ancient megafaunal mutualisms and extinctions as factors in plant domestication (2021, March 25) retrieved 30 May 2024 from <https://phys.org/news/2021-03-ancient-megafaunal-mutualisms-extinctions-factors.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.