


Biosensing with whispering-gallery mode lasers

March 16 2021



a, Single-cell monitoring with an intracellular microlaser. b, 3D arrangement of myofibrils around microbeads in neonatal cardiomyocytes (CMs). Cell nucleus (magenta) and microlaser (green). c, WGM spectrum of a microlaser and its shifting. d, Microlaser attached to the atrium of a zebrafish heart. e, Refractive index change between the resting phase, diastole, and peak contraction, systole, for 12 individual cells. f, Extracellular microlaser on top of an adult CM. Scale bar 30 μm . g, Trace of a spontaneously beating neonatal CM during administration of 500 nM nifedipine. Adapted with permission from Schubert M. et al. Monitoring contractility in cardiac tissue with cellular resolution using biointegrated microlasers. *Nature Photonics* 14, 452-458, (2020). Credit: Nikita Toropov, Gema Cabello, Mariana P. Serrano, Rithvik R. Gutha, Matías Rafti, Frank Vollmer

Whispering-gallery-mode (WGM) microresonators are opening up many new research directions that enable the detection of proteins, enzymes and DNA, down to single molecules. A new class of sensors makes use of active microresonators to drastically expand the functionality and range of WGM sensor applications, especially for biological and chemical assays. We review the most recent advances of WGM microlasers for biosensing and provide an outlook on exciting new research directions and emerging WGM sensor applications.

Label-free optical [sensors](#) based on optical whispering-gallery-mode (WGM) microresonators exhibit extraordinary sensitivity for detecting physical, chemical, and biological entities, even down to [single molecules](#). This advancement in label-free optical detection is made possible by application of the optical microresonator, i.e. a 100 um glass microspheres, as [optical cavity](#) to enhance the detection signal. Akin to a spherical micromirror, the WGM cavity reflects the light by near-total internal reflection and thereby creates multiple cavity passes that enhance the optical detection of analyte molecules interacting with the evanescent field.

In contrast to the 'cold' WGM microresonators, the emerging active WGM microlasers have the potential to significantly expand the number of possible applications of this class of sensors in biological and chemical sensing, and especially in in vivo sensing. The WGM microlasers can sense from within tissue, organisms and [single cells](#), and they can be used to improve upon the already impressive single-molecule detection limits of the 'cold'-cavity optoplasmonic WGM sensors.

Here, we review the most recent advances of WGM microlasers in biosensing. In contrast to the 'cold' cavity WGM sensors, the active WGM microresonators make use of gain media such as dye molecules

and [quantum dots](#) to compensate for optical loss and to achieve lasing of the WGM modes. Similar to other conventional lasers, lasing is observed from narrow spectral lines in the WGM emission spectra.

We review the main building blocks of WGM microlasers, recently demonstrated sensing mechanisms, the methods for integrating gain media in WGM sensors, and the prospects for active WGM sensors to become a useful technology in real-world applications. We review WGM [microlaser](#) sensing experiments at the molecular-level where lasing spectra are analyzed to study the binding of molecules, to sensing at the cellular level where microlasers are embedded into or integrated with single cells to enable novel in vivo sensing and single-cell tracking applications (see figure).

More information: Nikita Toropov et al, Review of biosensing with whispering-gallery mode lasers, *Light: Science & Applications* (2021).
[DOI: 10.1038/s41377-021-00471-3](https://doi.org/10.1038/s41377-021-00471-3)

Provided by Chinese Academy of Sciences

Citation: Biosensing with whispering-gallery mode lasers (2021, March 16) retrieved 13 March 2024 from <https://phys.org/news/2021-03-biosensing-whispering-gallery-mode-lasers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.