


Fluorescent nanodiamonds successfully injected into living cells

March 3 2021



SEM image of the FND used in this study. Credit: *Small* (2021). DOI: 10.1002/sml.202006421

As odd as it sounds, many scientists have attempted to place extremely small diamonds inside living cells. Why? Because nanodiamonds are consistently bright and can give us unique knowledge about the inner life of cells over a long time. Now physics researchers at Lund University in Sweden have succeeded in injecting a large number of nanodiamonds directly to the cell interior.

Diamonds are not only sought after for their beauty, but also for their uniquely luminescent properties, at least among scientists. Unlike other fluorescent materials, they do not bleach.


"We actually think of them as a dye. In addition, they are biocompatible," says Elke Hebisch, researcher at [solid state physics](#) at Lund University.

Together with Professor Christelle Prinz, she has "injected" fluorescent nano-sized diamonds into living cells.

As a researcher, having such a reporter from inside a cell has many advantages: gaining new knowledge about the cell, as well as monitoring what happens inside the cell over time.

"Especially the latter would be a great step forward, as it is currently possible to take snapshots of, for example, proteins in a cell, but difficult to follow changes over time," explains Elke Hebisch.

What would researchers want to know? It could be about separating [healthy cells](#) from diseased ones, targeting disease-causing proteins and other proteins within a specific cell, or monitoring variations in temperature and pH-levels. The knowledge gained could be pure basic research but can also be used to understand diseases and develop drugs.




Fast and high-efficiency intracellular delivery of FND via nanostraw-assisted injection, in comparison to low-yield FND delivery by incubation. a) Top panel: Schematics of the intracellular FND delivery via incubation. Middle and bottom panel: Confocal fluorescence microscopy images (xy, xz, and yz scans) of live A549 cells after 1 and 24 h of incubation in FND-containing medium. b) Top panel: Schematics of nanostraw-assisted FND injections. Bottom panel: Confocal fluorescence microscopy image of live cells on nanostraws imaged 5 min after FND delivery. In all images, the FND signal is shown in green-yellow-red, depending on the pixel intensity, and the cell nucleus signal is shown in blue. All images are 2D projections of the maximum pixel intensity obtained from 3D (XYZ) sample scans. For all images, the background noise corresponds to three counts. c) Quantification of the internalized FND signal: Integrated FND pixel intensity normalized to cell area (\pm S.E.M.) assessed for the

two FND delivery methods presented in (a) and (b). n.s.: $p > 0.05$; ****: $p \leq 0.0001$, two-sided Mann–Whitney–Wilcoxon U Test. Credit: *Small* (2021). DOI: 10.1002/smll.202006421

Other researchers have previously tried to do the same thing, but the diamonds were then taken care of by the cell's "cleaners," the so-called lysosomes, that quickly encapsulated the foreign substance.

"In that scenario, they are not useful since they are trapped in lysosomes and unable to interact with the cell components. Others have managed to get the [diamonds](#) into the cell one cell at a time, but that is far too time-consuming to become a realistic alternative," says Christelle Prinz.



Cell membrane recovery on nanostraws after the application of electroporation pulses. STED micrographs of the membrane of live A549 cells on top of nanostraws before, during, 5, 30, and 60 min after the application of low-voltage EP through the nanostraws. The membrane pores start to close after 30 min and are completely closed 60 min after switching off the EP. Credit:

The same technique could eventually be used to transport other molecules in order to alter cells or heal diseased [cells](#).

On a final note: Is using nanodiamonds expensive? No, Elke Hebisch

explains—the quantities needed are extremely small. They are bought in a bottle where they are suspended around in water, and cost the same as regular antibodies.

More information: Elke Hebisch et al. Nanostraw-Assisted Cellular Injection of Fluorescent Nanodiamonds via Direct Membrane Opening, *Small* (2021). [DOI: 10.1002/smll.202006421](https://doi.org/10.1002/smll.202006421)

Provided by Lund University

Citation: Fluorescent nanodiamonds successfully injected into living cells (2021, March 3) retrieved 21 May 2024 from <https://phys.org/news/2021-03-fluorescent-nanodiamonds-successfully-cells.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--