


High end of climate sensitivity in new climate models seen as less plausible

March 3 2021, by Morgan Kelly



The researchers found that models with lower climate sensitivity are more consistent with observed temperature differences, particularly between the northern and southern hemispheres. The graph shows changes in the annual global-mean surface temperature (a) and the temperature difference between the northern and southern hemispheres (b) from 1850 to 2000. The red line represent high climate-sensitivity models, while the blue line represents models with low climate sensitivity. The black line shows observed temperature fluctuations collected by NASA's Goddard Institute for Space Studies Surface Temperature Analysis project, which more closely follow the blue line when it comes to interhemispheric temperature. The gray backgrounds indicate years when the difference between the high and low climate-sensitivity models are significant. Credit: Chenggong Wang, Program in Atmospheric and Oceanic Sciences, Princeton University

A recent analysis of the latest generation of climate models—known as a CMIP6—provides a cautionary tale on interpreting climate simulations as scientists develop more sensitive and sophisticated projections of how the Earth will respond to increasing levels of carbon dioxide in the atmosphere.

Researchers at Princeton University and the University of Miami reported that newer models with a high "[climate](#) sensitivity"—meaning they predict much greater global warming from the same levels of atmospheric carbon dioxide as other models—do not provide a plausible scenario of Earth's future climate.

Those models overstate the global cooling effect that arises from interactions between clouds and aerosols and project that clouds will moderate [greenhouse gas](#)-induced warming—particularly in the [northern hemisphere](#)—much more than climate records show actually happens, the researchers reported in the journal *Geophysical Research Letters*.

Instead, the researchers found that models with lower climate sensitivity are more consistent with observed differences in temperature between the northern and southern hemispheres, and, thus, are more accurate depictions of projected climate change than the newer models. The study was supported by the Carbon Mitigation Initiative (CMI) based in Princeton's High Meadows Environmental Institute (HMEI).

These findings are potentially significant when it comes to climate-change policy, explained co-author Gabriel Vecchi, a Princeton professor of geosciences and the High Meadows Environmental Institute and principal investigator in CMI. Because models with higher climate sensitivity forecast greater warming from [greenhouse gas emissions](#), they also project more dire—and imminent—consequences such as more extreme sea-level rise and heat waves.

The high climate-sensitivity models forecast an increase in global average temperature from 2 to 6 degrees Celsius under current carbon dioxide levels. The current scientific consensus is that the increase must be kept under 2 degrees to avoid catastrophic effects. The 2016 Paris Agreement sets the threshold to 1.5 degrees Celsius.

"A higher climate sensitivity would obviously necessitate much more aggressive carbon mitigation," Vecchi said. "Society would need to reduce carbon emissions much more rapidly to meet the goals of the Paris Agreement and keep global warming below 2 degrees Celsius. Reducing the uncertainty in climate sensitivity helps us make a more reliable and accurate strategy to deal with climate change."

The researchers found that both the high and low climate-sensitivity models match global temperatures observed during the 20th century. The higher-sensitivity models, however, include a stronger cooling effect from aerosol-cloud interaction that offsets the greater warming due to greenhouse gases. Moreover, the models have aerosol emissions occurring primarily in the northern hemisphere, which is not consistent with observations.

"Our results remind us that we should be cautious about a [model](#) result, even if the models accurately represent past global warming," said first author Chenggong Wang, a Ph.D. candidate in Princeton's Program in Atmospheric and Oceanic Sciences. "We show that the global average hides important details about the patterns of temperature change."

In addition to the main findings, the study helps shed light on how clouds can moderate warming both in models and the real world at large and small scales.

"Clouds can amplify global warming and may cause [warming](#) to accelerate rapidly during the next century," said co-author Wenchang

Yang, an associate research scholar in geosciences at Princeton. "In short, improving our understanding and ability to correctly simulate clouds is really the key to more reliable predictions of the future."

Scientists at Princeton and other institutions have recently turned their focus to the effect that clouds have on climate change. Related research includes two papers by Amilcare Porporato, Princeton's Thomas J. Wu '94 Professor of Civil and Environmental Engineering and the High Meadows Environmental Institute and a member of the CMI leadership team, that reported on the future effect of heat-induced clouds on solar power and how climate models underestimate the cooling effect of the daily cloud cycle.

"Understanding how [clouds](#) modulate [climate change](#) is at the forefront of climate research," said co-author Brian Soden, a professor of atmospheric sciences at the University of Miami. "It is encouraging that, as this study shows, there are still many treasures we can exploit from historical climate observations that help refine the interpretations we get from global mean-temperature change."

The paper, "Compensation Between Cloud Feedback and Aerosol-Cloud Interaction in CMIP6 Models," was published in the Feb. 28 edition of *Geophysical Research Letters*

More information: Chenggong Wang et al, Compensation Between Cloud Feedback and Aerosol-Cloud Interaction in CMIP6 Models, *Geophysical Research Letters* (2021). [DOI: 10.1029/2020GL091024](https://doi.org/10.1029/2020GL091024)

Provided by Princeton University

Citation: High end of climate sensitivity in new climate models seen as less plausible (2021,

March 3) retrieved 24 April 2024 from <https://phys.org/news/2021-03-high-climate-sensitivity-plausible.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.