

'Lost' ocean nanoplastic might be getting trapped on coasts

March 10 2021



Credit: CC0 Public Domain

As plastic debris weathers in aquatic environments, it can shed tiny nanoplastics. Although scientists have a good understanding of how these particles form, they still don't have a good grasp of where all the

fragments end up. Now, researchers reporting in ACS' *Environmental Science & Technology* have shown experimentally that most nanoplastics in estuarine waters can clump, forming larger clusters that either settle or stick to solid objects, instead of floating on into the ocean.

There is a huge discrepancy between the millions of tons of plastic waste entering rivers and streams and the amount researchers have found in the oceans. As large pieces of plastic break apart into successively smaller fragments on their transit to the sea, some eventually wear down to nano-sized particles. Previous studies have shown that these nanoplastics congregate in well-mixed, stagnant salty [water](#). Yet, these results do not apply when the particles encounter dynamic changes in [salt content](#), such as estuaries, where rivers carrying freshwater meet tidal saltwater. So, Hervé Tabuteau, Julien Gigault and colleagues wanted to perform laboratory experiments with micro-sized chambers mimicking the conditions measured in an estuary to show how nanoplastics interact and aggregate in this type of environment.

To determine how nanoplastics move in estuarine waters, the team developed a lab-on-a-chip device. They introduced crushed 400-nm-wide polystyrene beads and freshwater into one side of the device, while injecting saltwater through another inlet. At the opposite end of the 1.7-cm-long device, the researchers collected the output. The team tested different flow rates, replicating the salt gradient and water movement they measured in an [estuary](#) on the French Caribbean island of Guadeloupe. Nanoplastic aggregates up to 10- μm wide were detected within the zone of highest salt concentration in the flow chamber, regardless of how fast the water was moving. At the highest flow rate, only 12% of the nanoplastics were collected in the outlets; the remaining particles either clumped together and sank in the flow chamber or formed floating aggregates that stuck to the chamber's sides. The researchers say their results show estuaries and other coastal environments may filter out nanoplastics before they can enter the

ocean.

More information: Zélie Venel et al. Environmental Fate Modeling of Nanoplastics in a Salinity Gradient Using a Lab-on-a-Chip: Where Does the Nanoscale Fraction of Plastic Debris Accumulate?, *Environmental Science & Technology* (2021). [DOI: 10.1021/acs.est.0c07545](https://doi.org/10.1021/acs.est.0c07545)

Provided by American Chemical Society

Citation: 'Lost' ocean nanoplastic might be getting trapped on coasts (2021, March 10) retrieved 11 May 2024 from <https://phys.org/news/2021-03-lost-ocean-nanoplastic-coasts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.