



Researchers notice pattern on surface of leaves, uncover new clue about plant evolution

March 29 2021



Plants in the genus *Callitriche* are short and have oval-shaped leaves about 1 cm in size. Amphibious *Callitriche palustris* (left, photographed in Matsuyama, Ehime Prefecture) often grows in small water channels in rice paddies. Terrestrial *Callitriche japonica* (right, photographed in Nagoya) can be found in parks or near shrines in urban areas. By studying these species, researchers at the University of Tokyo uncovered a pattern in how plants evolved their equivalent of lungs -- tiny pores on the surfaces of leaves called stomata. Credit: Hiroyuki

Koga, CC BY 4.0

A doctoral student has identified a long-overlooked pattern in how plants evolved their equivalent of lungs—tiny pores on the surfaces of leaves called stomata. Using specialized imaging techniques and a plant species not often found in laboratories, the researchers say this discovery reveals a key difference in the evolution of plants that live on land versus those that can grow in water.

"I felt this is really interesting, this was a big surprise to me. I remember well that after observation in the microscope room on the basement floor, I rushed up the stairs to tell Dr. Koga about my discovery," recalled first-year doctoral student Yuki Doll, studying in the University of Tokyo Graduate School of Science under the supervision of Assistant Professor Hiroyuki Koga.

"Of course, I and any scientist can see that the stomata are different, but it is easy for us to just ignore it, not sense any pattern. When I heard about Doll-kun's discovery, I was also very excited and discussed with him that we should delve into this subject," said Koga. (Kun is the Japanese honorific suffix attached to junior men's names.)

When stomata are open, carbon dioxide, oxygen and water vapor can move in and out of the leaf for photosynthesis and respiration. Artificially manipulating the number of stomata is one potential way to keep crops healthy in a changing climate.

The UTokyo team was studying multiple types of [plants](#) in the genus *Callitriche*, which includes both terrestrial and aquatic [species](#).

"*Callitriche* is an interesting but minor group of plants and we are the

only ones in the world using them for developmental biological research," said Koga.

Recalling his first experiences examining the plants, Doll said, "When I started to analyze stomata distribution patterns in aquatic *Callitriche*, I felt that the arrangement of the stomata are different than what I had been taught as an undergrad in the common lab species *Arabidopsis*. I had the impression that this strange pattern must be the case for all *Callitriche*, but I thought, that's OK, *Arabidopsis* and *Callitriche* are from very different evolutionary lineages, so it's natural for them to be different. Then I analyzed a terrestrial species of *Callitriche* and I saw it looked much more like *Arabidopsis*."

Specifically, Doll noticed that stomata and the cells surrounding them on the surface of aquatic plants' leaves were much more uniform than the variable cell sizes on the terrestrial plants' leaves.

This pattern that two evolutionarily closely related [plant species](#) had such different patterns of stomata development hinted at the possibility that their living conditions—on land or in water—might regulate stomatal development.

Koga and other lab members had previously perfected a method to visualize [gene activity](#) in every individual cell of intact, whole plant leaves. The technique of whole-mount fluorescence in situ hybridization is not new, but it is difficult and unusual to use those molecular biology tools without cutting a plant into ultrathin slices.

The images from terrestrial and aquatic *Callitriche* leaves confirmed that the plants used the same two genes to develop their stomata, but the genes were active at different times.

In almost all plants, the gene *SPEECHLESS* promotes growth and

division of a group of cells on the surfaces of newly forming leaves. Eventually, the gene MUTE becomes active in these cells and blocks SPEECHLESS, causing these cells to stop dividing and then differentiate to stomata. By binding artificial fluorescent tags to the two genes, researchers could see in single-cell resolution when SPEECHLESS is suppressed and MUTE becomes active.

In terrestrial *Callitriche*, researchers saw MUTE expressed in cells of all different ages. MUTE was much more uniformly expressed only in older [cells](#) of aquatic species, which seemed to skip the division stage and have a coordinated delay to wait until late in leaf development to activate MUTE.

Researchers suspect that [aquatic species](#) evolved to delay stomatal formation to wait and sense if this new leaf will be fully submerged or if it will be above the water line. Gas exchange is less efficient under water, so submerged leaves generally have fewer stomata than leaves in air.

The discovery is exciting for evolutionary biologists interested in the relationship between environmental pressures and evolutionary genetics, but is also relevant for the future of growing crops in changing or unpredictable environments.

"The usual assumption is that closely related species have similar stomata development patterns, but our key finding is that this is not the case," said Koga.

Instead, the researchers say their new results show that a species' living environment is the important evolutionary force selecting its stomata development pattern, not just the species' genetic ancestry.

By understanding the full genetic pathway that leads to flexible control

of SPEECHLESS and MUTE expression between species, scientists may be able to predict which evolutionary lineages of crops are more likely to optimize their [stomata](#) to grow in a changing climate.

More information: Yuki Doll et al., "The diversity of stomatal development regulation in *Callitriche* is related to the intrageneric diversity in lifestyles," *PNAS* (2021).

www.pnas.org/cgi/doi/10.1073/pnas.2026351118

Provided by University of Tokyo

Citation: Researchers notice pattern on surface of leaves, uncover new clue about plant evolution (2021, March 29) retrieved 11 May 2024 from <https://phys.org/news/2021-03-pattern-surface-uncover-clue-evolution.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.