


Pungency variable of Shishito sweet chili pepper: Genes and seeds

March 9 2021



Separation procedure for placental septa in 'Shishito' fruits. Each fruit was cut (a), and the placental septum was extracted (b). The placental septum was vertically separated (b), and one was used for capsaicinoid quantification and the other was used for qRT-PCR to measure the expression of the 18 genes involved in capsaicinoid biosynthesis. Credit: Copyright © 2021, Fumiya Kondo, Kenichi Matsushima et al., Shinshu University, under exclusive licence to Springer Nature Switzerland AG, part of Springer Nature


Chili peppers (*Capsicum spp.*) are an important spice and vegetable that supports food culture around the world, whose intensity of its pungent taste is determined by the content of capsaicinoids. However, the content of capsaicinoids varies depending on the variety and is known to fluctuate greatly depending on the cultivation environment. This can be a big problem in the production, processing and distribution of peppers where sweet varieties can be spicy and highly spicy varieties are just only mildly spicy. It is thought that changes in the expression of multiple genes involved in capsaicinoid biosynthesis are involved in such changes in pungent taste depending on the cultivation environment, but the mechanism is not clear.

In Japan, the main variety of pepper is the vegetable pepper shishito, which has almost no spiciness. However, sometimes spicy fruits are produced, which makes eating shishito like playing Russian roulette. People have known from experience that "fruits with a fewer number of seeds are spicy." Researchers set out to determine the number of seeds inside the shishito pepper and the intensity of [pungency](#) of the fruit, and the gene expression control mechanism that causes the fluctuation of the pungency of the shishito fruit.

Corresponding author, Associate Professor Kenichi Matsushima of the

Institute of Agriculture, Shinshu University states that the pungency of [chili peppers](#) with fewer seeds is more likely to fluctuate in pungency and that these pungency fluctuations are caused by the expression of multiple genes involved in capsaicinoid synthesis.

The research group investigated the pungent intensity of shishito fruits with a variety of number of seeds, with a tasting test and measurement of the capsaicinoid content. The placentation/partition (taiza/kakuheki) tissue in the fruit where capsaicinoids are synthesized was vertically divided into two equal parts, one was measured for pungent intensity and the other for the expression of genes involved in capsaicinoid synthesis. As a result, the spiciness of the shishito fruit with a large number of seeds was very weak, whereas the spiciness of the fruits with a small number of seeds ranged from weak to strong.




Pungency traits and the number of seeds in commercially available 'Shishito' fruits according to sensory evaluations. a The number of seeds in non-pungent fruits and pungent fruits. In each boxplot, the maximum and minimum values (as whiskers), first and third quartiles (as a boundary box), the median (as a horizontal line in the box), and the mean (as a cross-mark) are indicated. P

The group investigated the relationship between the intensity of pungency and the degree of gene function within the same [fruit](#). As a result, among the genes involved in capsaicinoid synthesis, the [expression levels](#) of 10 genes showed a positive correlation with the pungent intensity, and it was found that the higher the expression level of these genes, the stronger the pungent taste. Therefore, it was clarified that in shishito [peppers](#), the activation of these 10 genes increases the amount of capsaicinoids synthesized and enhances the pungent taste.

There have been many studies investigating how much the pungent intensity (capsaicinoid content) of peppers changes depending on the cultivation conditions, but few studies have investigated gene expression. In this study, the pungent fluctuation phenomenon of chili pepper was investigated at the gene expression level. The group clarified the relationship between the two by using a unique experimental method of simultaneously investigating pungent intensity and gene expression by using the placenta and septum of shishito in two equal parts.

In addition to the results of this research, this novel method can also be considered to be an important outcome for investigating the pungent fluctuation phenomenon in the future. Associate Professor Matsushima hopes to utilize the knowledge and information on the expression level of genes that have been found to be significantly involved in pungent fluctuations for future pungency breeding of chili peppers. It was revealed in this study that the pungent [intensity](#) of pepper and the expression level of multiple genes involved in capsaicinoid synthesis are closely related.



Capsaicinoid biosynthesis pathways and the involvement of the 18 analyzed genes in the present study. Letters in each circle indicate the gene name and the ten co-expressed genes whose expression levels had positive correlations ($P < 0.05$, $P < 0.01$, $P < 0.001$).

Based on this result, if the expression level of these genes can be suppressed, it may be possible to grow vegetable varieties that are less likely to fluctuate in pungency. They are aiming for a variety of shishito that does not produce irregularly of spicy fruits, while retaining the unique flavor, which is different from the peppers that are not spicy at all. Therefore, the results of this research will be applied to pepper breeding, and the ultimate goal is to establish breeding technology for these [genes](#) and their functions.

More information: Fumiya Kondo et al, The pungent-variable sweet chili pepper 'Shishito' (*Capsicum annuum*) provides insights regarding the relationship between pungency, the number of seeds, and gene expression involving capsaicinoid biosynthesis, *Molecular Genetics and Genomics* (2021). [DOI: 10.1007/s00438-021-01763-4](https://doi.org/10.1007/s00438-021-01763-4)

Provided by Shinshu University

Citation: Pungency variable of Shishito sweet chili pepper: Genes and seeds (2021, March 9)
retrieved 6 August 2024 from

<https://phys.org/news/2021-03-pungency-variable-shishito-sweet-chili.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.