

## Found in space: Complex carbon-based molecules

March 18 2021, by Sarah McDonnell



The Taurus Molecular Cloud, which contains the cold starless core TMC-1, is a dark streak on the sky near the Pleiades cluster as seen from Charlottesville, VA. Credit: Brett A. McGuire, Copyright 2018

Much of the carbon in space is believed to exist in the form of large molecules called polycyclic aromatic hydrocarbons (PAHs). Since the



1980s, circumstantial evidence has indicated that these molecules are abundant in space, but they have not been directly observed.

Now, a team of researchers led by MIT Assistant Professor Brett McGuire has identified two distinctive PAHs in a patch of space called the Taurus Molecular Cloud (TMC-1). PAHs were believed to form efficiently only at high temperatures—on Earth, they occur as byproducts of burning fossil fuels, and they're also found in char marks on grilled food. But the interstellar cloud where the research team observed them has not yet started forming stars, and the temperature is about 10 degrees above absolute zero.

This discovery suggests that these molecules can form at much lower temperatures than expected, and it may lead scientists to rethink their assumptions about the role of PAH chemistry in the formation of stars and planets, the researchers say.

"What makes the detection so important is that not only have we confirmed a hypothesis that has been 30 years in the making, but now we can look at all of the other molecules in this one source and ask how they are reacting to form the PAHs we're seeing, how the PAHs we're seeing may react with other things to possibly form larger molecules, and what implications that may have for our understanding of the role of very large carbon molecules in forming planets and stars," says McGuire, who is a senior author of the new study.

Michael McCarthy, associate director of the Harvard-Smithsonian Center for Astrophysics, is another senior author of the study, which appears today in *Science*. The research team also includes scientists from several other institutions, including the University of Virginia, the National Radio Astronomy Observatory, and NASA's Goddard Space Flight Center.



## **Distinctive signals**

Starting in the 1980s, astronomers have used telescopes to detect infrared signals that suggested the presence of aromatic molecules, which are molecules that typically include one or more carbon rings. About 10 to 25 percent of the carbon in space is believed to be found in PAHs, which contain at least two <u>carbon rings</u>, but the infrared signals weren't distinct enough to identify specific molecules.

"That means that we can't dig into the detailed chemical mechanisms for how these are formed, how they react with one another or other molecules, how they're destroyed, and the whole cycle of carbon throughout the process of forming stars and planets and eventually life," McGuire says.

Although radio astronomy has been a workhorse of molecular discovery in space since the 1960s, radio telescopes powerful enough to detect these large molecules have only been around for a little over a decade. These telescopes can pick up molecules' rotational spectra, which are distinctive patterns of light that molecules give off as they tumble through space. Researchers can then try to match patterns observed in space with patterns that they have seen from those same molecules in laboratories on Earth.





The 100-m Green Bank Telescope located in Green Bank, WV. Credit: Brett A. McGuire, Copyright 2018

"Once you have that pattern match, you know there is no other molecule in existence that could be giving off that exact spectrum. And, the intensity of the lines and the relative strength of the different pieces of the pattern tells you something about how much of the molecule there is, and how warm or cold the molecule is," McGuire says.

McGuire and his colleagues have been studying TMC-1 for several years because previous observations have revealed it to be rich in complex carbon molecules. A few years ago, one member of the research team observed hints that the cloud contain benzonitrile—a six-carbon ring attached to a nitrile (carbon-nitrogen) group.



The researchers then used the Green Bank Telescope, the world's largest steerable radio telescope, to confirm the presence of benzonitrile. In their data, they also found signatures of two other molecules—the PAHs reported in this study. Those molecules, called 1-cyanonaphthalene and 2-cyanonaphthalene, consist of two benzene rings fused together, with a nitrile group attached to one ring.

"Detecting these molecules is a major leap forward in astrochemistry. We are beginning to connect the dots between small molecules—like benzonitrile—that have been known to exist in space, to the monolithic PAHs that are so important in astrophysics," says Kelvin Lee, an MIT postdoc who is one of the authors of the study.

Finding these molecules in the cold, starless TMC-1 suggests that PAHs are not just the byproducts of dying stars, but may be assembled from smaller molecules.

"In the place where we found them, there is no star, so either they're being built up in place or they are the leftovers of a dead star," McGuire says. "We think that it's probably a combination of the two—the evidence suggests that it is neither one pathway nor the other exclusively. That's new and interesting because there really hadn't been any observational evidence for this bottom-up pathway before."





In a series of nine papers, scientists from the GOTHAM--Green Bank Telescope Observations of TMC-1: Hunting Aromatic Molecules--project described the detection of more than a dozen polycyclic aromatic hydrocarbons in the Taurus Molecular Cloud, or TMC-1. These complex molecules, never before detected in the interstellar medium, are allowing scientists to better understand the formation of stars, planets, and other bodies in space. In this artist's conception, some of the detected molecules include, from left to right: 1-cyanonaphthalene, 1-cyanocyclopentadiene, HC11N, 2-cyanonaphthalene, vinylcyanoacetylene, 2-cyanocyclopentadiene, benzonitrile, trans-(E)-cyanovinylacetylene, HC4NC, and propargylcyanide, among others. Credit: M. Weiss / Center for Astrophysics I Harvard & Smithsonian

## **Carbon chemistry**



Carbon plays a critical role in the formation of planets, so the suggestion that PAHs might be present even in starless, cold regions of space, may prompt scientists to rethink their theories of what chemicals are available during planet formation, McGuire says. As PAHs react with other molecules, they may start to form interstellar dust grains, which are the seeds of asteroids and planets.

"We need to entirely rethink our models of how the chemistry is evolving, starting from these starless cores, to include the fact that they are forming these large aromatic molecules," he says.

McGuire and his colleagues now plan to further investigate how these PAHs formed, and what kinds of reactions they may undergo in <u>space</u>. They also plan to continue scanning TMC-1 with the powerful Green Bank Telescope. Once they have those observations from the <u>interstellar</u> cloud, the researchers can try to match up the signatures they find with data that they generate on Earth by putting two molecules into a reactor and blasting them with kilovolts of electricity, breaking them into bits and letting them recombine. This could result in hundreds of different molecules, many of which have never been seen on Earth.

"We need to continue to see what <u>molecules</u> are present in this interstellar source, because the more we know about the inventory, the more we can start trying to connect the pieces of this reaction web," McGuire says.

**More information:** B.A. McGuire el al., "Detection of two interstellar polycyclic aromatic hydrocarbons via spectral matched filtering," *Science* (2021). <u>science.sciencemag.org/cgi/doi ... 1126/science.abb7535</u>

Ci Xue et al. Detection of Interstellar HC4NC and an Investigation of Isocyanopolyyne Chemistry under TMC-1 Conditions, *The Astrophysical Journal* (2020). <u>iopscience.iop.org/article/10. ... 847/2041-8213/aba631</u>



Brett A. McGuire et al. Early Science from GOTHAM: Project Overview, Methods, and the Detection of Interstellar Propargyl Cyanide (HCCCH2CN) in TMC-1, *The Astrophysical Journal* (2020). iopscience.iop.org/article/10. ... 847/2041-8213/aba632

Andrew M. Burkhardt et al. Ubiquitous aromatic carbon chemistry at the earliest stages of star formation, *Nature Astronomy* (2021). DOI: 10.1038/s41550-020-01253-4

Michael C. McCarthy et al. Interstellar detection of the highly polar fivemembered ring cyanocyclopentadiene, *Nature Astronomy* (2020). DOI: 10.1038/s41550-020-01213-y

Provided by Massachusetts Institute of Technology

Citation: Found in space: Complex carbon-based molecules (2021, March 18) retrieved 17 July 2024 from <u>https://phys.org/news/2021-03-space-complex-carbon-based-molecules.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.