

Rating tornado warnings charts a path to improve forecasts

March 2 2021, by Hannah Hickey



A funnel cloud from a tornado in Kansas on May 24, 2016, inside the United States' so-called "Tornado Alley." Credit: Lane Pearman/Flickr

The United States experiences more tornadoes than any other country, with a season that peaks in spring or summer depending on the region.

Tornadoes are often deadly, especially in places where buildings can't withstand high winds.

Accurate advanced warnings can save lives. A study from the University of Washington and the National Oceanic and Atmospheric Administration describes a new way to rate and possibly improve [tornado warnings](#). It finds that nighttime twisters, summer [tornadoes](#) and smaller events remain the biggest challenges for the forecasting community.

"This new method lets us measure how forecast skill is improving, decreasing or staying the same in different situations," said Alex Anderson-Frey, a UW assistant professor of atmospheric sciences. "The tornado forecasting community needs to know what we're doing best at, and where we can focus training and research in the future."

She is lead author of the paper published online in December in the *Bulletin of the American Meteorological Society*.

Though the southern and central U.S. see the most tornadoes, every state can experience twisters. Scientific understanding of tornadoes is biased toward populated places, Anderson-Frey said, where people are more likely to observe and report the events.

"As population density increases in different areas, including outside the U.S., I think we're getting more of an idea of the range of environments in which tornadoes can actually form," Anderson-Frey said.

The paper develops a new method to rate the skill of a tornado [warning](#) based on the difficulty of the environment. It then evaluates thousands of tornadoes and associated warnings over the continental United States between 2003 and 2017.

The NOAA-funded study finds that nighttime tornadoes have a lower probability of detection and a higher false-alarm rate than the [environmental conditions](#) would suggest. Summertime tornadoes, occurring in June, July or August, also are more likely to evade warning.



Tornadoes are most common in the southeastern U.S., but these small-scale atmospheric events can form in many places. This EF-5 tornado was photographed as it approached Elie, Manitoba, late in the afternoon of June 22, 2007. Credit: Justin1569/Wikimedia

The nighttime events may be harder to forecast because the absence of

daytime warming makes the conditions less favorable, and because there were fewer eyewitness reports, Anderson-Frey said. Summer events may be more difficult because summer has more relatively weak tornadoes that occur in marginal environments, meaning on the edge of conditions that produce a tornado.

Larger events—those rated 2 or above on the enhanced Fujita scale—actually generated better warnings than expected for the conditions. The results can inform how research, training or observational technology could improve future tornado warnings.

"The forecasting community is not just looking at the big, photogenic situations that will crop up in the Great Plains. We're looking at tornadoes in regions where vulnerability is high, including in regions that don't normally get tornadoes, where by definition the vulnerability is high," Anderson-Frey said.

"There's a real effort in the forecasting research community to bring in the human element—being able to identify where we can do the most good."

Although tornado forecasts and warnings are improving overall, so are some types of risk. Populations are growing and moving into new, remote environments. Mobile or manufactured homes without anchored foundations are less able to withstand high winds.

"What really excites me about this work is the opportunity to look at performance by how difficult the warning situation was," said co-author Harold Brooks at NOAA's National Severe Storm Laboratory in Norman, Oklahoma. "We have the chance to measure improvement through the years taking into account that some situations and years may be harder or easier forecasts."

Anderson-Frey moved from Oklahoma to join the UW faculty in 2019. In related research, she is now analyzing data for past tornadoes to determine the environmental conditions that can produce events in unexpected places, like the 2018 tornado that struck Port Orchard in Washington state.

"I'm working on applying a machine-learning technique to studying what prototypical tornadic environments look like in different parts of the United States," she said.

More information: Alexandra K. Anderson-Frey et al, Compared to What? Establishing Environmental Baselines for Tornado Warning Skill, *Bulletin of the American Meteorological Society* (2020). [DOI: 10.1175/BAMS-D-19-0310.1](https://doi.org/10.1175/BAMS-D-19-0310.1)

Provided by University of Washington

Citation: Rating tornado warnings charts a path to improve forecasts (2021, March 2) retrieved 18 April 2024 from <https://phys.org/news/2021-03-tornado-path.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--