


Drones provide bird's eye view of how turbulent tidal flows affect seabird foraging habits

April 27 2021



A picture of the SeaGen tower in Strangford Lough, Northern Ireland, taken from a drone. Credit: Alex Nimmo-Smith, University of Plymouth

The foraging behavior of seabirds is dramatically affected by turbulence caused by natural coastal features and manmade ocean structures, new research has shown.

In a first-of-its-kind study, scientists from the UK and Germany used drones to provide a synchronized bird's eye view of what seabirds see and how their behavior changes depending on the movement of tidal flows beneath them.

The research focused on the wake of a tidal turbine structure set in a tidal channel—Strangford Lough in Northern Ireland—that has previously been identified as a [foraging](#) hotspot for terns.

Through a combination of drone tracking and advanced statistical modeling, it showed that terns were more likely to actively forage over vortices (swirling patches of water).

However, eruptions of upwelling water (boils) ahead of the terns' [flight path](#) prompted them to stay on course as they approached.

Writing in the Royal's Society flagship biological research journal, *Proceedings of the Royal Society B*, the researchers say their findings offer a never-before-seen insight into how tidal turbulence can impact foraging behaviors.

They also say it potentially gives them the ability to predict how species might respond to [environmental changes](#) such as the increased future development of ocean renewable energy sites and [climate change](#).



A tern carries away food caught during a successful foraging dive. Credit: Lilian Lieber, Queen's University Belfast

The study was conducted by researchers from Queen's University Belfast and the University of Plymouth (UK), and Bielefeld University (Germany).

Dr. Lilian Lieber, Bryden Centre Research Fellow at Queen's and the study's lead investigator, said: "Our research highlights the importance of identifying changes in local flow conditions due to ocean energy structures which can change the occurrence, scale and intensity of localized turbulence in the water. Through a fantastic interdisciplinary collaboration, we were able to track prevalent flow features and seabirds on thus far unobtainable scales, shedding new light on tern foraging

associations with turbulence. We found that terns were more likely to actively forage over vortices, while conspicuous upwellings provided a strong physical cue even at some distance, leading them to investigate such features. This research can help us predict seabird responses to coastal change."

Co-investigator Professor Roland Langrock, Professor in Statistics and Data Analysis at Bielefeld, said: "It is extremely exciting that we now have these incredibly detailed animal movement data, which allows us to investigate behavioral processes at effectively arbitrarily fine scales of animal decision-making. While it presented some new statistical challenges, the interdisciplinary nature of our project presents a valuable contribution to the emerging field of high-throughput movement ecology."

Co-investigator Dr. Alex Nimmo-Smith, Associate Professor in Marine Physics in Plymouth, led the computational development of automatically and reliably tracking the terns using machine learning as well as mapping the underlying turbulent features.

He added: "The drone provided a real bird's eye view, allowing us to track the highly localized foraging behavior of the terns and the close association they have with particular flow features. Upwelling boils and swirling vortices, characteristic of strong tidal flows, can bring potential prey items (such as small fish) to the water surface and trap them there. Therefore, these physical processes provide foraging opportunities for the terns."

More information: A bird's eye view on turbulence: Seabird foraging associations with evolving surface flow features, *Proceedings of the Royal Society B*, rspb.royalsocietypublishing.org/doi/10.1098/rspb.2021.0592

Provided by University of Plymouth

Citation: Drones provide bird's eye view of how turbulent tidal flows affect seabird foraging habits (2021, April 27) retrieved 24 April 2024 from <https://phys.org/news/2021-04-drones-bird-eye-view-turbulent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.