

Study: Drought-breaking rains more rare, erratic in US West

April 6 2021, by Matthew Brown



The time between rainfalls has become longer and the rains occurred more erratically in the western U.S. during the last 50 years. Credit: Joel Biederman, Agricultural Research Service

Rainstorms grew more erratic and droughts much longer across most of the U.S. West over the past half-century as climate change warmed the

planet, according to a sweeping government study released Tuesday that concludes the situation is worsening.

The most dramatic changes were recorded in the desert Southwest, where the average dry period between rainstorms grew from about 30 days in the 1970s to 45 days between storms now, said Joel Biederman, a research hydrologist with the U.S. Department of Agriculture Southwest Watershed Research Center in Tucson, Arizona.

The consequences of the intense dry periods that pummeled areas of the West in recent years were severe—more intense and dangerous wildfires, parched croplands and not enough vegetation to support livestock and wildlife. And the problem appears to be accelerating, with rainstorms becoming increasingly unpredictable, and more areas showing longer intervals between storms since the turn of the century compared to prior decades, the study concludes.

The study comes with almost two-thirds of the contiguous U.S. beset by abnormally dry conditions. Warm temperatures forecast for the next several months could make it the [worst spring drought](#) in almost a decade, affecting roughly [74 million people](#) across the U.S., the National Oceanic and Atmospheric Administration said.

Water use cutbacks, damaged wheat crops, more fires and lower reservoirs in California and the Southwest are possible, weather service and agriculture officials have warned. Climate scientists are calling what's happening in the West a continuation of a "megadrought" that started in 1999.

While previous research documented a decline in total rainfall for much of the West, the work by Biederman and colleagues put more focus on when that rain occurs. That has significant implications for how much water is available for agriculture and plants such as grasses that have

shallow roots and need a steadier supply of moisture than large trees.

"Once the growing season starts, the total amount of rainfall is important. But if it comes in just a few large storms, with really long dry periods in between, that can have really detrimental consequences," study co-author Biederman said in an interview.

The total amount of rain in a year doesn't matter to plants—especially if rains come mostly in heavy bursts with large run-off—but consistent moisture is what keeps them alive, said UCLA meteorologist Daniel Swain, who writes a weather blog about the West and was not part of the study.

The new [findings were published in the journal *Geophysical Research Letters*](#). Researchers led by University of Arizona climate scientist Fangyue Zhang compiled daily readings going back to 1976 from 337 weather stations across the western U.S. and analyzed rainfall and drought data to identify the changing patterns.

Other parts of the region that saw longer and more variable droughts included the southwest Rocky Mountains, the Colorado Plateau and the Central Plains.

The rainfall study is in line with data that shows climate change already is affecting the planet.

"Climate models project that the American Southwest is very likely to experience more frequent and more severe droughts," said William Anderegg, a University of Utah biologist and climate scientist. "This study and other recent work demonstrates that this dry down has already begun."

The weather station data that was used in the study represents "the gold

standard' for an accurate understanding of changes being driven by climate change, said Christopher Field, an earth systems scientist and director of the Stanford Woods Institute for the Environment.

Park Williams—who studies changes in water, wildfires and climate at UCLA—cautioned more work was needed to see if the rainfall trends since the 1970s reflect a longer-term natural cycle or are tied to human-caused warming. Regardless, the combination of longer dry spells and warmer temperatures almost certainly adds to increased wildfire size, he said.

Northwestern states were largely spared from the accelerating cycles of drought. The researchers observed higher annual rainfall totals and shorter drought intervals in Washington, Oregon, Idaho and portions of Montana, Wyoming and the Dakotas.

That's consistent with predicted alterations in weather patterns driven by climate change in which the jet stream that brings moisture from the Pacific Ocean shifts northward, they said.

More information: *Geophysical Research Letters* (2021). [DOI: 10.1029/2020GL092293](https://doi.org/10.1029/2020GL092293)

© 2021 The Associated Press. All rights reserved. This material may not be published, broadcast, rewritten or redistributed without permission.

Citation: Study: Drought-breaking rains more rare, erratic in US West (2021, April 6) retrieved 18 April 2024 from <https://phys.org/news/2021-04-droughts-longer-rainfall-erratic-years.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.