


# Insect evolution was more complex than previously assumed

April 21 2021

---



University of Cologne scientists found out that the Toll signaling pathway is important not only for innate immune response, but also for axis formation in various insects. Credit: Roth/Pechmann

Certain signaling proteins, which are responsible for the development of innate immune function in almost all animals are also required for the

formation of the dorsal-ventral (back-belly) axis in insect embryos. A new study by researchers from the University of Cologne's Institute of Zoology suggests that the relevance of these signaling proteins for insect axis formation has increased independently several times during evolution. For example, the research team found similar evolutionary patterns in the Mediterranean field cricket as in the fruit fly *Drosophila*, although the two insects are only very distantly related and previous observations suggested different evolutionary patterns. The new findings show that the evolution of axis formation in insects was actually much more complex than previously thought. The study has been published in *eLife*.

Signaling proteins play an important role in the early development of embryos. They are secreted by [animal cells](#) to influence the formation of other cells. The primary function of the so-called Toll signaling pathway is in the defense against pathogens (innate immune response). In insects, it is also involved in the division of the insect body along the dorsal-ventral body axis. Since the immune function has been found in almost all animals, but the axis formation function has only been found in insects, scientists wondered about the evolutionary history of this new role.

Moreover, depending on the [insect species](#), the significance of Toll for developmental processes differs. While axis formation in the fruit fly and flour beetle depend substantially on Toll, representatives of distantly [related species](#), such as the wasp *Nasonia* and the milkweed bug *Oncopeltus*, rely more heavily on other signaling pathways. 'Surprisingly, we found that the Toll signaling pathway plays a significant role in an insect that is separated by almost 400 million years from the species we studied so far,' said Professor Dr. Siegfried Roth from the Institute of Zoology. 'The new study suggests that there might be several instances in which Toll independently acquired important functions in insect [axis](#) formation. For future studies, this means that our system allows us to

explore mechanisms of parallel evolution.'

**More information:** Matthias Pechmann et al, Striking parallels between dorsoventral patterning in *Drosophila* and *Gryllus* reveal a complex evolutionary history behind a model gene regulatory network, *eLife* (2021). [DOI: 10.7554/eLife.68287](https://doi.org/10.7554/eLife.68287)

Provided by University of Cologne

Citation: Insect evolution was more complex than previously assumed (2021, April 21) retrieved 23 May 2024 from <https://phys.org/news/2021-04-insect-evolution-complex-previously-assumed.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.