

New material for catholytes and anolytes in organic redox flow batteries

April 6 2021



Credit: CC0 Public Domain



Researchers from Skoltech and their collaborators have designed, synthesized and evaluated new compounds that can serve as catholytes and anolytes for organic redox flow batteries, bringing this promising technology closer to large-scale implementation. The two papers were published in the *Journal of Materials Chemistry A* and *Chemical Communications*.

Energy storage is a crucial component of a greener energy system of the future based on <u>renewable sources</u>; batteries have to be complemented with wind and solar farms and have to be scalable, safe, and flexible in their design and lifetime. Redox flow batteries (RFBs) are all those things, yet one major barrier to commercialization has been their low specific capacity. So, a lot of research effort is focused on developing better battery components to overcome this hurdle.

"The main advantage of <u>redox-flow batteries</u> is scalability—the capacity of the battery is limited only by the volume of the electrolyte, so it is the ideal construction for large-scale <u>energy storage</u>. Nowadays we are working with organic redox-active materials solubilized in <u>organic</u> <u>solvents</u> (non-aqueous organic RFBs). The main advantages for nonaqueous organic RFB are high cell voltage (up to 5V, versus around 1.6 V for water-based systems), a huge variety of organic redox-active molecules which cold be applied, and potential operability at low temperatures, without any concern for freezing below 0 degrees C. As such this work offers considerable advances for the development of RFBs of this type," Skoltech Ph.D. student Elena Romadina, the first author of both papers, explains.

In the two papers, Elena Romadina and her colleagues describe highly promising catholyte and anolyte materials for RFBs—triarylamine-based materials and a phenazine derivative, respectively. The seven highly soluble redox-active triarylamine-based compounds were designed, synthesized and tested for solubility and electrochemical properties, with



one of them, singled out as the most promising candidate for further studies. The authors emphasize that the developed compounds exhibited almost unlimited solubility in polar organic solvents such as acetonitrile, which makes them promising for high-capacity RFBs. In the other study, a phenazine derivative with oligomeric ethylene glycol ether substituents was synthesized in a two-step process and showed solid performance as a RFB anolyte.

"A non-aqueous organic redox flow battery designated as phenazinebased anolyte and most promising triarylamine-based catholyte exhibited a high cell voltage of 2.3 V, high capacity, >95% coulombic efficiency and good charge-discharge cycling stability during the 50 cycles," the authors write in the *ChemComm* paper.

"As a result of our work, we presented a novel class of compounds that could be used in RFBs. Previously, poly-triarylamines were investigated as a cathode material for metal-ion cells, but this class of compounds was not investigated in redox-flow batteries. Thus, a new and very promising core structure was opened up for us and other scientists. Triarylamines have a stable and fully reversible redox potential, and could be easily modified, providing different redox potentials and physical properties. Moreover, we found that triarylamines-based compounds could retain their electrochemical properties even in the presence of water in organic solvent, which lowered the requirements for solvent preparation and cost," Romadina adds.

"We are in fact looking at both ends of the battery in an effort to raise operational cell voltage and to prevent other degradation of catholytes and anolytes. To make organic RFBs commercially viable, we also need research in areas such as low-cost scalable synthesis of highly soluble redox-active molecules; the development of high-performance membranes that are good ionic conductors, but inhibit cross-over of anolytes and catholytes upon charge and discharge; and the scaling of



larger cell and stack level device configurations to enable grid scale energy storage," Professor Keith Stevenson, Skoltech provost and coauthor of the papers, says.

More information: Elena I. Romadina et al. New highly soluble triarylamine-based materials as promising catholytes for redox flow batteries, *Journal of Materials Chemistry A* (2021). DOI: 10.1039/D0TA11860E

Elena I. Romadina et al. New phenazine based anolyte material for high voltage organic redox flow batteries, *Chemical Communications* (2021). DOI: 10.1039/D0CC07951K

Provided by Skolkovo Institute of Science and Technology

Citation: New material for catholytes and anolytes in organic redox flow batteries (2021, April 6) retrieved 2 May 2024 from <u>https://phys.org/news/2021-04-material-catholytes-anolytes-redox-batteries.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.