

# Study reveals extent of human impact on the world's plant-life

April 29 2021

---



Santo Antao (Cabo Verde). Credit: Sandra Nogue

Research has shed new light on the impact of humans on Earth's biodiversity. The findings suggest that the rate of change in an ecosystem's plant-life increases significantly during the years following

human settlement, with the most dramatic changes occurring in locations settled in the last 1500 years.

An international research team studied fossilized pollen dating back 5000 years, extracted from sediments on 27 [islands](#). By analyzing the fossils they were able to build up an understanding of the composition of each island's vegetation and how it changed from the oldest to the most recent pollen samples.

The study was led by Dr. Sandra Nogué, Lecturer in Palaeoenvironmental Science at the University of Southampton, UK and Professor Manuel Steinbauer from the University of Bayreuth, Germany and University of Bergen, Norway. Ph.D. student Dr. Alvaro Castilla-Beltrán was also a member of the Southampton team.

Dr. Nogué said, "Islands provide the ideal environment to measure [human impact](#) as most were settled in the past 3000 years when climates were similar to today's conditions. Knowing when the settlers arrived on an island means that scientists can study how the composition of its ecosystem changed in the years before and after."

The results, published in *Science*, showed a consistent pattern on 24 of the islands where human arrival accelerated the turnover of vegetation by, on average, a factor of eleven. The most rapid changes occurred in islands that were settled more recently—such as the Galápagos, first inhabited in the 16th Century. Islands where humans arrived more than 1500 years ago, such as Fiji and New Caledonia, saw a slower rate of change.





Teide Fieldwork - Canary Islands. Credit: José María Fernández Palacios

"This difference in change could mean that the islands populated earlier were more resilient to human arrival but it is more likely that the land-use practices, technology and introduced species brought in by the later settlers were more transformative than those of the earlier settlers," explained Dr. Nogué.

The trends were observed across a range of geographic locations and climates, with islands such as Iceland producing similar results to Tenerife and other tropical and temperate islands.



Ecosystem change can also be driven by a number of natural factors such as earthquakes, [volcanic eruptions](#), [extreme weather](#) and changing sea levels, however the researchers have found that disturbance caused by humans surpasses all of these events and the change is often irreversible. They therefore advise that conservation strategies must account for the long-term impact of humans and the degree to which ecological changes today differ from prehuman times.



Field Work in Cabo Verde. Credit: Sandra Nogue

"Whilst it is unrealistic to expect ecosystems to return to their pre-settlement conditions, our findings may help to inform targeted

[restoration efforts](#) and provide greater understanding into the islands' responsiveness to change," concludes Dr. Nogué.

**More information:** "The human dimension of biodiversity changes on islands" *Science* (2021). [science.sciencemag.org/cgi/doi ... 1126/science.abd6706](https://science.sciencemag.org/cgi/doi/10.1126/science.abd6706)

Provided by University of Southampton

Citation: Study reveals extent of human impact on the world's plant-life (2021, April 29) retrieved 16 August 2024 from <https://phys.org/news/2021-04-reveals-extent-human-impact-world.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--