

Young male fruit flies make females fight each other more

April 21 2021, by Dr Ellie Bath



Image 1: Two female fruit flies are standing on a food cap (which contains food that they eat and lay eggs in). This is where the majority of fighting happens – you can see the flies have their legs touching, so they are probably fencing in this image. Each fly is marked with a different colour of paint to aid identification. Credit: University of Oxford

Mating changes female behavior across a wide range of animals, with these changes induced by components of the male ejaculate, such as sperm and seminal fluid proteins. However, males can vary significantly

in their ejaculates, due to factors such as age, mating history, or feeding status. This male variation may therefore lead to variation in the strength of responses males can stimulate in females.

Using the fruit fly, *Drosophila melanogaster*, we tested whether age, mating history, and feeding status shape an important, but understudied, post-mating response—increased female-female aggression.

We found that [females](#) mated to old males fought less than females mated to [young males](#). Females mated to old, sexually active males fought even less than those mated to males who were merely old, but there was no effect of male starvation status on mating-induced female aggression.

Male condition can therefore influence how females interact with each other—who you mate with changes your interactions with members of the same sex!

"Male condition influences female post [mating](#) aggression and feeding in *Drosophila*" is published in *Functional Ecology*.

Could this happen in other species?

We know that other species (including humans!) have proteins in the seminal fluid that [males](#) transfer to females during sex. Various of these proteins have effects on female physiology and behavior, but no one knows if these affect aggression (in humans or any other species).




Image 2: This figure shows the setup we used for contests between females, which consisted of a circular arena with a food cap set into the middle. This food cap contained regular fly food medium, with a drop of yeast paste in the middle to act as a valuable, restricted resource. Credit: University of Oxford

We also know that male age (in flies and humans) results in reduced fertility and can have serious effects on their offspring. Although it is a long leap from flies to humans, could who you mate with influence your interactions with other females?

Many reproductive molecules and important bodily functions are conserved across the animal kingdom from flies to humans, so it is possible that what we found in flies here might be hinting to a common phenomenon across the tree of life. We need more studies to understand if this is the case!

More information: Eleanor Bath et al. Male condition influences female post mating aggression and feeding in *Drosophila*, *Functional*

Ecology (2021). [DOI: 10.1111/1365-2435.13791](https://doi.org/10.1111/1365-2435.13791)

Provided by University of Oxford

Citation: Young male fruit flies make females fight each other more (2021, April 21) retrieved 19 June 2024 from <https://phys.org/news/2021-04-young-male-fruit-flies-females.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.