


# **Aquaculture turns biodiversity into uniformity along the coast of China**

May 26 2021

---



Bivalves were sampled at 21 sites along the Chinese coast from the city of Dongliadao in the tropical south, to the mudflats of Yalu Jiang. Credit: He-bo Peng (NIOZ/RUG)

Fishery and aquaculture have given rise to an enormous uniformity in the diversity of bivalves along the more than 18,000 kilometer long Chinese coast, biologist He-Bo Peng and colleagues report in this month's issue of *Diversity and Distributions*.

## **Climate zones**

Peng and colleagues sampled bivalves at 21 sites along the Chinese coast from the city of Dongliadao in the tropical south, to the mudflats of Yalu Jiang, more than 2000 km further north and ice-covered for several months in winter. "At 19 out of these 21 sites, commercially exploited species dominated", Peng saw. "In the naturally occurring species, we still recognized the natural gradient with highest diversity in the tropics and lowest diversity in the north. However, the same commercially exploited species were found in all regions, regardless the climatic conditions. Also, the commercial species dominated almost all mudflat communities."

## **Shrimp fisheries**

One of the [dominant species](#) nowadays is *Potamocorbula laevis*. Peng says, "At some locations, this bivalve represented 95% of the biomass." This is not necessarily due to the culturing of this bivalve. "The intensive fishing for shrimps has almost eradicated a natural enemy of the small fragile juveniles of this bivalve. Possibly, this lack of predators has now

favored *P. laevis*."

## **Blessing in disguise**

The eradication of the south to north biodiversity gradient and the dominance of commercially exploited bivalve [species](#) may be a blessing in disguise for [migratory birds](#) along the Chinese coast. Peng: "At sites where there's a lot of *p. laevis* now, migratory birds like red and great knots find a lot of food as they refuel during their migrations between Australia and the tundra areas of northeast Russia. But the more general point raised by this study is the immensity of the role of human activities in determining coastline biodiversity. Where once there was a strong climate gradient in biodiversity, aquaculture and fisheries have favored uniformity now."

## **Continuous sampling**

"This extensive research along more than 18,000 km of Chinese coast builds on the Dutch tradition of studying mudflat ecosystems", Theunis Piersma, researcher at NIOZ and promotor of Ph.D.-student Peng as professor of global flyway ecology at the University of Groningen, says. "Sampling programs like SIBES in the Dutch Wadden Sea taught us the value of widespread and continuous sampling. In this study He-Bo Peng once again shows the value of documenting benthos at very large spatial scales, with sampling strategies developed in the Wadden Sea."

**More information:** He-Bo Peng et al, Mollusc aquaculture homogenizes intertidal soft-sediment communities along the 18,400 km long coastline of China, *Diversity and Distributions* (2021). [DOI: 10.1111/ddi.13302](https://doi.org/10.1111/ddi.13302)

Provided by Royal Netherlands Institute for Sea Research

Citation: Aquaculture turns biodiversity into uniformity along the coast of China (2021, May 26)  
retrieved 11 May 2024 from <https://phys.org/news/2021-05-aquaculture-biodiversity-uniformity-coast-china.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.