


Discovered: Cell component that turns the staphylococcal bacterium deadly

May 25 2021



Digitally colorized scanning electron microscopic image of *Staphylococcus aureus* bacteria (mustard-colored) enmeshed within a human white blood cell (red-colored). Credit: National Institute of Allergy and Infectious Diseases (NIAID)

The bacterium *Staphylococcus epidermidis* is a usually harmless microbe found on the skin and in the noses of humans. Yet some strains of this species can cause infections—in catheters, artificial joints, heart valves and in the bloodstream, and are difficult to treat. These bacteria are often resistant to a particularly effective antibiotic, methicillin, and are among the most feared germs in hospitals. How these usually harmless skin microbes become deadly pathogens has been unclear up to now.

An international research team has now discovered what distinguishes peaceful *S. epidermidis* microorganisms from the many dangerous invaders. The scientists have identified a new gene cluster that enables the more aggressive bacteria to produce additional structures in their cell walls. This morphological alteration allows the staphylococci to attach more easily to [human cells](#) forming the blood vessels, a process via which they can persist in the bloodstream to become pathogens. These new [cell wall](#) structures may also allow the spread of methicillin resistance, by transferring it, for example, from *Staphylococcus epidermidis* to its more dangerous relative *Staphylococcus aureus*.

The study was carried out under the direction of researchers of the Cluster of Excellence "Controlling Microbes to Fight Infections" (CMFI) of the University of Tübingen and the German Center for Infection Research (DZIF) in cooperation with universities in Copenhagen, Hamburg, Shanghai and Hanover and the German Center for Lung Research (DZL) in Borstel. The results are being published in the journal *Nature Microbiology*.

Set apart by structure

A considerable portion of the cell walls of Staphylococci—like other [gram-positive bacteria](#)—is made up of teichoic acids. These chain-like polymers cover the bacterial surface. Their chemical structures vary

according to species. "During our examination, we determined that many pathogenic strains of *S. epidermidis* have an additional gene cluster that contains information for the synthesis of wall teichoic acids that are actually typical of *S. aureus*," says researcher Dr. Xin Du of the Cluster of Excellence of the CMFI and DZIF. She adds that experiments have shown *S. epidermidis* bacteria with only species-specific teichoic acids in their walls are not very invasive, colonizing the surfaces of the skin and mucous membranes. If the wall teichoic acids for *S. aureus* are also present, Xin Du explains, they are unable to attach effectively to those surfaces. Instead, they are more successful in penetrating the tissues of their human host. "At some point, a few *S. epidermidis* clones took on the corresponding genes from *S. aureus* and became threatening pathogens as a result," says Professor Andreas Peschel of the Cluster of Excellence CMFI and of the DZIF.

It's long been known that bacteria can share genetic material through gene transfer. Bacteriophages—viruses that infect bacteria—carry out the transfer. Mostly, this takes place within one species and requires similar surface structures to which the bacteriophages bind. "Differing cell wall structures normally prevent gene transfer between *S. epidermidis* and *S. aureus*. But in *S. epidermidis* strains that can also produce the wall teichoic acids of *S. aureus*, that type of gene transfer suddenly becomes possible between different species," explains Peschel. That would explain how *S. epidermidis* could transfer methicillin resistance to even more threatening—and then methicillin-resistant—*S. aureus*; he says that more investigation is still needed. The new findings are an important step, says Peschel, towards developing better treatments or vaccinations against dangerous pathogens such as *S. epidermidis* ST 23, which has been known for 15 years and belongs to the group of HA-MRSE (healthcare-associated methicillin-resistant *S. epidermidis*).

More information: Xin Du et al, *Staphylococcus epidermidis* clones express *Staphylococcus aureus*-type wall teichoic acid to shift from a

commensal to pathogen lifestyle, *Nature Microbiology* (2021). [DOI: 10.1038/s41564-021-00913-z](https://doi.org/10.1038/s41564-021-00913-z)

Provided by University of Tübingen

Citation: Discovered: Cell component that turns the staphylococcal bacterium deadly (2021, May 25) retrieved 2 April 2024 from <https://phys.org/news/2021-05-cell-component-staphylococcal-bacterium-deadly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.