

## **Convective transport explains 'missing' ice near the tropical tropopause**

May 11 2021, by David Shultzdavid Shultz



The orange-red glow in this image is Earth's troposphere, the lowest layer of the atmosphere. The brown transitional layer is the tropopause. In a new study, spaceborne lidar shows more ice than expected leaving the tropopause over the tropics. Credit: <u>NASA Earth Observatory</u>, <u>CC BY 2.0</u>



The lowest level of the atmosphere, the troposphere, contains almost all of Earth's weather. In the stratosphere above, moisture drops to almost zero. The boundary that separates these two layers—the tropopause—is defined as the point at which water ceases to cool as altitude increases. In the tropics, the tropopause is exceptionally cold and usually occurs at a higher altitude (around 17 kilometers, or about 10 miles) than in polar regions. The atmospheric conditions at these heights play an important role in global climate, as they determine the limits of cloud formation and water vapor dynamics.

In a new study, Bolot and Fueglistaler use spaceborne lidar to study ice fluxes near the tropical tropopause. They used NASA's Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) instrument to calculate how much ice is falling out of the tropopause layer, known as sedimentation. Significantly, more ice is found than ambient moisture can account for at these altitudes.

Ice near the tropical <u>tropopause</u> can come from two sources: advection of cirrus clouds and <u>deep convection</u>. The first can be thought of simply as large, icy clouds being lifted upward by background motions, whereas deep convection represents upward movement along a thermal gradient—a buoyancy of water/ice particles resulting from warmer, less dense material rising. By using data from ERA5 (a fifth-generation European Centre for Medium-Range Weather Forecasts atmospheric reanalysis), the researchers show that advection alone is insufficient to account for the missing ice below 17 kilometers and that the remainder must be made up by deep convection.

**More information:** Maximilien Bolot et al. Tropical Water Fluxes Dominated by Deep Convection Up to Near Tropopause Levels, *Geophysical Research Letters* (2021). DOI: 10.1029/2020GL091471



## Provided by Eos

Citation: Convective transport explains 'missing' ice near the tropical tropopause (2021, May 11) retrieved 28 June 2024 from <u>https://phys.org/news/2021-05-convective-ice-tropical-tropopause.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.