

Coral fights back against crown of thorns starfish

May 5 2021, by Ivy Shih



A small juvenile crown of thorns starfish (approx. 15 mm) retreating after being stung by coral polyps. Credit: Dione Deaker

Coral can fight back against attacking juvenile crown of thorns



starfish—using stinging cells to injure and even kill, showing that coral are not as passive as people may think.

Coral are not completely defenseless against attacking juvenile <u>crown</u> of thorns <u>starfish</u> and can fight back to inflict at times lethal damage, new research has found.

This occurs during a period of the crown of thorns starfish life cycle, where small juveniles shift from a vegetarian diet of algae to coral prey. But this change in diet makes the juveniles more vulnerable to attack by coral.

Population outbreaks of adult crown of thorns starfish, alongside <u>coral</u> <u>bleaching</u> is one of the greatest threats to tropical reef habitats.

Video footage shows when the tube feet (small tube-like projections on the underside of a starfish's arm used for movement) of juvenile crown of thorns starfish reaches out to touch the coral, the entire arm curls back to avoid the corals' defensive stinging cells. To protect themselves, coral polyps have stinging cells in their sweeper tentacles and outer tissue called nematocysts, that are also used to capture food.

This encounter damages the arms of juvenile crown of thorn starfish, delaying their growth into adulthood. Researchers also saw a 10 percent fatality rate among the juvenile crown of thorns starfish they studied. However, most juveniles that survived arm damage were able to regenerate partially lost arms.





An injured juvenile crown of thorns starfish. Credit: Dione Deaker

The research, published in *Marine Ecology Progress Series*, was led by Dione Deaker, a Ph.D. student at the University of Sydney, and her supervisor Professor Maria Byrne. The marine scientists say that this is the first study of injury and regeneration in juvenile crown of thorn starfish following damage caused by natural enemies.



The researchers emphasize the results give a fascinating insight into coral behavior but the behavior is not enough to protect it from other threats such as human-caused climate change, overfishing and water pollution.

Ms Deaker says the period when young crown of thorns starfish shift from a vegetarian diet to eating coral, which is an animal, is a critical one. This is because young crown of thorns starfish who survive have the potential to contribute to population outbreaks that could devastate tropical reefs and coral.

<u>Previous research</u> led by Ms Deaker and Professor Byrne has shown juvenile starfish can survive on algae for more than six years when they were previously thought to change diets at four months old, lying in wait until there is an abundance of coral.

Caught on tape





A juvenile crown of thorns starfish with ultimately fatal damage to the arms and central disk. Credit: Dione Deaker

Marine biologists have reported seeing injured juvenile starfish and have suggested that it may be been caused by predators.

"However, seeing it caused by coral came as a complete surprise," said



Ms Deaker.

"This shows that the coral use stinging cells as protection to strike back in an attempt to give itself a fighting chance against attacking coral predators."

In the study, Ms Deaker and Professor Byrne, along with colleagues at the national Marine Science Center, Coffs Harbor, monitored the condition, growth and survival of 37 juvenile crown of thorns in isolation away from potential predators and reared them on a diet of coral prey for over 3 months.

They found coral stings caused injuries that severely reduced the arm length of the starfish by up to 83 percent.

37.8 percent of juveniles were damaged by coral and four juveniles (10.8 percent) with severe injuries did not recover and died.





A juvenile crown of thorns starfish with arm regeneration after injury. Credit: Dione Deaker

The sting attacks from the coral also delayed the growth of juveniles, extending the time they need to maintain a vegetarian diet.

The young starfish had a reflex response to being stung when they encountered coral. Their arms recoiled and twisted when their tube feet



came into contact with the coral polyps.

"Sometimes the juveniles never recovered and died, but in most cases injured juveniles recovered and can regenerate their arms in about 4 months," said Ms Deaker.

"Despite being prey of crown of thorns starfish, coral can potentially influence the survival of juveniles and the appearance of a population outbreak on a reef by delaying their transition into an adult that can reproduce."

Armed with these observations, the study shows that coral are a risky food choice for young crown of thorns starfish.

Although coral injury was able to slow down the growth of the juvenile starfish, their ability to regenerate shows the resilience of this reef predator as a highly prolific species.

Professor Byrne said: "The importance of this study in showing the disconnect between size and age of the juveniles reinforces how challenging it is to understand the dynamics of adult population replenishment."

More information: Dione J. Deaker et al. The hidden army: corallivorous crown-of-thorns seastars can spend years as herbivorous juveniles, *Biology Letters* (2020). DOI: 10.1098/rsbl.2019.0849

DJ Deaker et al. Coral defences: the perilous transition of juvenile crown-of-thorns starfish to corallivory, *Marine Ecology Progress Series* (2021). DOI: 10.3354/meps13660



Provided by University of Sydney

Citation: Coral fights back against crown of thorns starfish (2021, May 5) retrieved 15 May 2024 from <u>https://phys.org/news/2021-05-coral-crown-thorns-starfish.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.