

Mangroves and seagrasses absorb microplastics

May 4 2021



Many animals live in seagrass beds and are thus exposed to microplastic. Credit: Troels Lange/SDU.



Mangroves and seagrasses grow in many places along the coasts of the world, and these 'blue forests' constitute an important environment for a large number of animals. Here, juvenile fish can hide until they are big enough to take care of themselves; crabs and mussels live on the bottom; and birds come to feed on the plants.

However, the plant-covered <u>coastal zones</u> do not only attract <u>animals</u> but also microplastics, a new study shows.

"The denser the vegetation, the more plastic is captured," says Professor and expert in coastal ecology, Marianne Holmer, from the University of Southern Denmark.

She is concerned about how the accumulated microplastics affect animal and plant life.

"We know from other studies that animals can ingest microplastics and that this may affect their organism."

Animals ingest microplastics with the food they seek in the blue forests. They may suffocate, die of starvation, or the small plastic particles can get stuck different places in the body and do damage.

Another problem with microplastics is that they may be covered with microorganisms, environmental toxins or other health hazardous/disease-promoting substances that are transferred to the animal or plant that absorbs the microplastics.

"When microplastics are concentrated in an ecosystem, the animals are exposed to very high concentrations," Marianne Holmer explains.

She points out that microplastics concentrated in, for example, a seagrass bed are impossible to remove again.



The study is based on examinations of three <u>coastal areas</u> in China, where mangroves, Japanese eelgrass (Z. japonica) and the paddle weed Halophila ovalis grow. All samples taken in blue forests had more microplastics than samples from control sites without vegetation.

The concentrations were up to 17.6 times higher, and they were highest in the <u>mangrove</u> forest. The concentrations were up to 4.1 times higher in the seagrass beds.

Mangrove trees probably capture more microplastics, as the capture of particles is greater in <u>mangrove forests</u> than in seagrass beds.

Researchers also believe that microplastics bind in these ecosystems in the same way as carbon; the particles are captured between leaves and roots, and the microplastics are buried in the seabed.

"Carbon capture binds carbon dioxide in the seabed, and the blue forests are really good at that, but it's worrying if the same thing happens to microplastics," says Marianne Holmer.

Although the study was conducted along Chinese coasts, it may be relevant to similar ecosystems in the rest of the world, including Denmark, where eelgrass beds are widespread.

"It's my expectation that we will also find higher concentrations of microplastics in Danish and global seagrasses," she says.

The study was conducted in collaboration with colleagues from the Zhejiang University in China, among others, and is published in the journal *Environmental Science and Technology*.

More information: Yuzhou Huang et al, New Insights into the Microplastic Enrichment in the Blue Carbon Ecosystem: Evidence from



Seagrass Meadows and Mangrove Forests in Coastal South China Sea, *Environmental Science & Technology* (2021). DOI: <u>10.1021/acs.est.0c07289</u>

Provided by University of Southern Denmark

Citation: Mangroves and seagrasses absorb microplastics (2021, May 4) retrieved 16 August 2024 from <u>https://phys.org/news/2021-05-mangroves-seagrasses-absorb-microplastics.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.