

Physicists extract proton mass radius from experimental data

May 12 2021



Fig. 1. Vector meson near threshold photoproduction process on the proton target. Credit: KOU Wei

Researchers have recently extracted the proton mass radius from experimental data. A research group at the Institute of Modern Physics (IMP) of the Chinese Academy of Sciences (CAS) presented an analysis of the proton mass radius in *Physical Review D* on May 11. The proton



mass radius is determined to be 0.67 ± 0.03 femtometers, which is obviously smaller than the charge radius of the proton.

In the Standard Model, the <u>proton</u> is a composite particle made of quarks and gluons and it has a non-zero size. The radius of the proton is a global and fundamental property of the proton. It is related to the color confinement radius—a property governed by quantum chromodynamics (QCD).

The radius of the proton is approximately 100,000 times smaller than that of the atom, and the sizes of the quark and gluon are several orders smaller than the proton radius. Scientists use various distributions to describe the shape of the proton, such as charge distribution and mass distribution.

The charge radius of the proton has been precisely measured by scientists via Lamb shift of the muonic hydrogen or the high energy electron-proton elastic scattering, with the average value of 0.8409 ± 0.0004 femtometers provided by the Particle Data Group. Nevertheless, knowledge of proton gravitational properties such as proton mass radius has still been very limited.

"According to recent theoretical studies by Dmitri Kharzeev, the proton mass radius is related to the scalar gravitational form factor of the proton," said Dr. Wang Rong, first author of the paper. By investigating the vector meson photoproduction data for omega, phi and J/psi from the SAPHIR (Spectrometer Arrangement for PHoton Induced Reactions) experiment at Bonn University, the LEPS (Laser Electron Photons) experiment at SPring-8 facility, and the GlueX experiment at Jefferson Lab, the researchers determined the scalar gravitational form factor and the proton mass radius.





Fig. 2. Proton mass radius extracted from experimental data. Credit: Physical Review D

Meanwhile, Prof. Dmitri Kharzeev, a <u>theoretical physicist</u> at Stony Brook University, obtained a similar result by using GlueX J/psi data. The proton mass radius was estimated to be 0.55 ± 0.03 femtometers.

"Both results might be the first-ever values of the proton mass radius with experimental evidence," said Wang. "The determination of the proton mass radius will improve our understanding of the origins of proton mass and the color confinement mechanism of strong interaction."



A lot of questions still remain. "The smaller mass radius implies that the <u>mass distribution</u> is significantly different from the charge distribution of the proton," said Prof. Chen Xurong, a researcher at IMP.

Scientists are now trying to get a clearer picture of the proton <u>mass</u> radius and the proton structure. The GlueX experiment at Jefferson Lab will provide more data in the near future. Even more exciting, future electron-ion colliders both in the United States and in China will provide Upsilon vector meson electroproduction data for researchers to better understand these questions.

More information: Rong Wang et al, Extraction of the proton mass radius from the vector meson photoproductions near thresholds, *Physical Review D* (2021). DOI: 10.1103/PhysRevD.103.L091501

Provided by Chinese Academy of Sciences

Citation: Physicists extract proton mass radius from experimental data (2021, May 12) retrieved 28 June 2024 from https://phys.org/news/2021-05-physicists-proton-mass-radius-experimental.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.