

## **Solar development: Super bloom or super bloot of the set of the s**

May 3 2021



The rare Barstow woolly sunflower was more sensitive to solar development impacts than its common relative, the woolly daisy in a study by UC Davis and UC Santa. Credit: Karen Tanner



Throughout the history of the West, human actions have often rushed the desert—and their actions backfired. In the 1920s, the Colorado River Compact notoriously overallocated water still used today by several western states because water measurements were taken during a wet period.

More currently, operators of the massive Ivanpah Solar Electric Generating System in the Mojave Desert are spending around \$45 million on desert tortoise mitigation after initial numbers of the endangered animals were undercounted before its construction.

A study published in the journal *Ecological Applications* from the University of California, Davis, and UC Santa Cruz warns against another potential desert timing mismatch amid the race against <u>climate</u> <u>change</u> and toward rapid renewable energy development.

"Our study suggests that <u>green energy</u> and <u>species</u> conservation goals may come into conflict in California's Mojave Desert, which supports nearly 500 rare plant species as well as a rapidly expanding solar industry," said lead author Karen Tanner, who conducted the work as a Ph.D. student at UC Santa Cruz under a grant led by UC Davis assistant professor Rebecca R. Hernandez.

Tanner spent seven years teasing out the demography of two native desert flowers—the rare Barstow woolly sunflower (E. mohavense) and the common Wallace's woolly daisy (E. wallacei), comparing their performance both in the open and under experimental solar panels. The authors wondered, how would desert-adapted plants respond to panels that block light and rainfall? Would rare species respond differently than <u>common species</u> to these changes?

These aren't easy questions to unearth. At one point, Tanner glued tiny seeds to individual toothpicks to gather emergence data. At another, she



scoured the desert floor on hands and knees to count emerging seedlings of the rare sunflower—about the size of a thumbnail at maturity.



A common Wallace's woolly daisy grows in the Mojave Desert. Common wildflowers appear to be less vulnerable than rare wildflowers to desert solar developments, a study from UC Davis and UC Santa Cruz found. Credit: Karen Tanner

## Super bloom surprises

Such painstaking commitment is one reason no previous studies have



modeled species' responses to photovoltaics at the population level. It takes time and overcoming tricky logistical and mathematical challenges to model little-known species interactions in the evasive desert. What is nowhere in sight one year, may thrive the next.

That element of surprise is what makes "super blooms" so special and so captivating. Those bursts of wildflowers blanket expanses of desert landscapes after especially wet years and are believed to be critical to the long-term persistence of desert annual populations.

The study found that solar panel effects on plant response were strongly influenced by weather and physical features of the landscape. During the 2017 super bloom, panel shade negatively affected population growth of the rare species, but had little effect on its common relative.

The study suggests that <u>rare species</u> may be more sensitive to solar development impacts than common species. It highlights the potential for solar panel effects to vary among species, as well as over space and time.





Widflowers blanket the desert near the study site in the Mojave Desert. Credit: Karen Tanner

## A question of time

The study provides an example of the importance of taking the necessary time to understand an ecosystem before irrevocably changing it.

"The <u>desert</u>—and many other biomes—don't respond on our timescales," said Hernandez, co-director of the <u>Wild Energy Initiative</u> through the UC Davis John Muir Institute. "If we want to understand them, we need to study them on the timescales they operate. Otherwise, it is like taking



a photo of a moving train and calling it a shipping container. Racing to build renewable energy in places that have already been skinned of their biology makes sense—let's not wait to put solar on existing rooftops. But in natural environments, we need to listen and observe first."

**More information:** Karen E. Tanner et al, Microhabitats associated with solar energy development alter demography of two desert annuals, *Ecological Applications* (2021). <u>DOI: 10.1002/eap.2349</u>

Provided by UC Davis

Citation: Solar development: Super bloom or super bust for desert species? (2021, May 3) retrieved 26 April 2024 from <u>https://phys.org/news/2021-05-solar-super-bloom-species.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.