

Growing sweet corn at higher densities doesn't increase root lodging risk

May 10 2021, by Lauren Quinn



Credit: Pixabay/CC0 Public Domain

Sweet corn growers and processors could be bringing in more profits by exploiting natural density tolerance traits in certain hybrids. That's according to 2019 research from USDA Agricultural Research Service



(ARS) and University of Illinois scientists.

But since <u>root systems</u> get smaller as plant <u>density</u> goes up, some in the industry are concerned about the risk of root lodging with greater sweet corn density. New research says those concerns are unjustified.

"Root lodging can certainly be a problem for sweet corn, but not because of plant density. What really matters is the specific hybrid and the environment, those major rainfall and wind events that set up conditions for root structural failure," says Marty Williams, USDA-ARS ecologist, affiliate professor in the Department of Crop Sciences at Illinois, and author on a new study in *Crop Science*.

Williams and his co-authors used multiple approaches to understand the effects of planting density on root lodging in sweet corn. First, they planted sweet corn hybrid DMC 21-84—a density-tolerant type that happens to be one of the most widely grown commercial hybrids—at five densities in experimental plots on U of I farms. When plants were at the tasseling stage, the researchers simulated a natural lodging event by flooding the field and knocking the corn over with a two-by-four.

"We mimicked a root lodging event with pure brute force," Williams says.

When corn is flattened in the process of root lodging, it can, depending on the growth stage, right itself through the plant's natural inclination to grow toward the light. As corn grows back upward from its repose, the base of the stem often carries a curved reminder of its lodging legacy, known as a gooseneck.

Williams and his team found most plants recovered to a near vertical position, albeit with a bit of goosenecking, within a few days of the artificial lodging event. They also measured yield metrics, and found no



statistical difference in sweet corn yield between lodged and non-lodged plants.

But a two-by-four and brute force can't replace nature.

"Testing a crop's root lodging potential experimentally is inherently difficult. Simply put, a natural root lodging event may not happen in any given field experiment. As such, phenotyping crops for root lodging often employs the use of artificially created lodging events, like the ones we created," Williams says. "While artificially created root lodging events are helpful, used alone they often fail to capture a broad range of environments in which the crop is grown."

That's why the research team leveraged data from on-farm sweet corn trials across Illinois, Minnesota, and Wisconsin. Between 2013 and 2017, the team evaluated economic optimum density for DMC 21-84 and 10 other hybrids under real-world farm management. And during that time, natural lodging events happened to occur in six of the 30 fields.

"We happened to take notes on lodging severity. It wasn't our specific focus at the time, but we figured, let's go ahead and score this and maybe we'll use it later. Turns out it was great that we collected root lodging data, because we were able to use it for this study," Williams says.

Those fortuitous notes showed that when sweet <u>corn</u> density increased from the current standard to the economic optimum density, typically a few thousand more plants per acre, there was absolutely no difference in the severity of lodging.

"What excited me about this work is that we combined an experiment where we created lodging artificially and looked at the response, and then we also tapped into this network of naturally occurring events. Together, they told a pretty convincing story," Williams says. "Simply



put, root lodging potential should not keep us from using <u>plant density</u> tolerant hybrids and growing them at their correct density. I mean, there's always the possibility that root lodging could occur, but it's not going to be due to planting a few more <u>plants</u>."

More information: Martin Williams et al, Economic optimum plant density of sweet corn does not increase root lodging incidence, *Crop Science* (2021). DOI: 10.1002/csc2.20546

Provided by University of Illinois at Urbana-Champaign

Citation: Growing sweet corn at higher densities doesn't increase root lodging risk (2021, May 10) retrieved 1 May 2024 from <u>https://phys.org/news/2021-05-sweet-corn-higher-densities-doesnt.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.