

# Research team traces ten million years of climate history in the Arabian Peninsula

May 18 2021

---



Zagros Mountains on the edge of Mesopotamia (Iran). Credit: Madelaine Böhme

The formation of deserts on the Arabian Peninsula had a decisive impact

on the migration and evolution of large mammals and our human ancestors over millions of years. That is the conclusion of a new study by an international research team led by Professor Madelaine Böhme of the Senckenberg Centre for Human Evolution and Palaeoenvironment at the University of Tübingen. The researchers reconstructed the climate history of the northern Arabian Peninsula from 12.5 to 2.5 million years ago using aridity-sensitive data obtained from rocks from Mesopotamia. This yielded new clues about the causes of animal migrations. The results have been published in the journal *Communications Earth & Environment*.

The evolution of today's African savannah fauna took place in relative isolation over the past five million years. This has been known for some time, as has the fact that the ancestors of many savannah animals such as rhinos, giraffes, hyenas and big cats came from Eurasia. However, what led the animals to make this large-scale change of location between the continents was unclear until now.

## Climate data stored in rock

The northern Arabian Peninsula is the gateway to Africa. Today, it includes [desert areas](#) such as the Syrian Desert, the Negev Desert in Israel and the Saudi Nefud Desert, as well as less dry steppes and semi-deserts in Mesopotamia, the largest part of which is in present-day Iraq. The research team studied the 2.6-kilometer-thick rock layers at the foot of the Zagros Mountains in Iran, on the edge of Mesopotamia, using chemical, physical and geological methods. The researchers found evidence for four brief phases of desertification in Mesopotamia, each lasting only a few tens of thousands of years. These phases, 8.75 million, 7.78 million, 7.5 million and 6.25 million years ago, were each interrupted by periods of wetter climate. "5.6 million years ago, coinciding with the temporary drying-out of the Mediterranean Sea, Mesopotamia experienced an extreme drought that lasted for 2.3 million

years," says Madelaine Böhme. This exceptionally long-lasting period of [desert](#) climate—referred to by Böhme's team as NADX (Neogene Arabian Desert climaX) – was only ended by [global warming](#) 3.3 million years ago, she says.



Researchers examined 12 million years of climate history of the northern Arabian Peninsula on 2.6 kilometer-thick rock layers. Credit: University of Tübingen

## Phases of migration and isolation



"Contrary to what we had expected, these desert phases on the Arabian Peninsula did not coincide with those in the African Sahara," Böhme reports. Desertification in the Sahara was causally linked to polar ice formations, she says, while that on the Arabian Peninsula and in Mesopotamia was linked to a low water table in the Caspian Sea, according to the findings. "The reciprocal emergence and decay of deserts in the Sahara in northern Africa, on the one hand, and on the Arabian Peninsula in western Asia, on the other, is like a kind of swing, a desert swing," Böhme says.

The research team assumes that these changing and initially short-lived deserts formed in Mesopotamia were the push factor behind the spread of mammals from Eurasia to Africa. By contrast, during the subsequent extremely long-lasting desert phase NADX, the African continent was cut off from migration and exchange with Eurasia for 2.3 million years, she said. "During this time, the Eurasian arrivals gave rise to today's African savanna fauna, and the australopithecids evolved, our [human ancestors](#)," Böhme explains. During the global warm period 3.3 million years ago, the deserts in both continents shrank, ending Africa's isolation, and a mutual exchange developed between the faunas of Africa and Eurasia. The first dogs, pigs and sheep appeared in Africa, and the ancestors of the mammoth and the Asian elephant migrated to Eurasia.

## Explanation for two phenomena

"Climatological explanations for two central phenomena emerge for the first time from our study," Böhme sums up. On the one hand, these underpin Böhme's Out-Of-Europe hypothesis, according to which the ancestors of African apes and humans evolved in Europe, but dispersed south six to seven million years ago, so that their further evolution took place in Africa. On the other hand, this could explain why the evolution of the African savannah fauna, including human ancestors, took place during a long period of isolation.

**More information:** Madelaine Böhme et al. Neogene hyperaridity in Arabia drove the directions of mammalian dispersal between Africa and Eurasia, *Communications Earth & Environment* (2021). [DOI: 10.1038/s43247-021-00158-y](https://doi.org/10.1038/s43247-021-00158-y)

Provided by University of Tübingen

Citation: Research team traces ten million years of climate history in the Arabian Peninsula (2021, May 18) retrieved 28 April 2024 from <https://phys.org/news/2021-05-team-ten-million-years-climate.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--