


Lead levels in urban soil are declining but hotspots persist

May 27 2021



Urban soil processes tend to reduce and dissipate soil lead contamination from paint and gasoline, which may be relevant for public health approaches. Credit: *Environmental Science & Technology* (2021). DOI: 10.1021/acs.est.1c00546

Decades after federal bans ended widespread use of lead in paint and gasoline, some urban soils still contain levels of the highly toxic metal that exceed federal safety guidelines for children, a Duke University study finds.

To conduct their study, the researchers analyzed and mapped [soil](#) lead concentrations along 25 miles of streets in Durham, N.C., a [city](#) of about 270,000 people. They found that while soil lead levels have generally decreased since the 1970s, they have decreased much less near residential foundations than along streets.

The researchers collected [soil samples](#) near foundations of houses built before 1978. Samples within a meter of the older homes averaged 649 milligrams (mg) of lead per kilogram (kg) of soil, more than three times the average level detected near streets, which was 150 mg/kg.

EPA guidelines say exposure to soil lead concentrations above 400 mg/kg is associated with potential long-term health risks to children, including possible damage to the brain and nervous system, slowed growth and development, learning and behavior problems and hearing and speech problems.

"Urban soil processes are driving lead concentrations down over time, but it's alarming that lead levels in some locations—typically older, poorer neighborhoods—still far exceed safe levels decades after [leaded gasoline](#) and lead paint were phased out," said lead author Anna Wade, a postdoctoral researcher at the U.S. Environmental Protection Agency and a 2020 Ph.D. graduate of Duke's Nicholas School of the Environment.

The Duke team shared its findings with Durham public health groups and plans to conduct similar mapping studies in five or six other cities nationwide.

Determining where contamination risks persist, and why contamination is decreasing at different rates in different locations, is essential for mitigating those risks, she said. However, many cities lack the resources to conduct the regular city-wide sampling needed to obtain that data. "Our study had to go all the way back to the 1970s to find comparable data for portions of the 35-square-kilometer area we sampled," Wade said.

Daniel D. Richter, professor of soils at Duke, said, "There's been a lot of interest in mitigating lead exposure in cities, but most is focused on reducing risks within the home. Our study reminds us about the outdoor environment where exposure risks also exist."

Wade, Richter and their team published their peer-reviewed, open-access study May 21 in *Environmental Science & Technology*.

Curbside soil lead levels have dropped over time as the result of human and [natural causes](#), the study suggests. Those causes include accelerated erosion and stormwater runoff, which has carried away some contaminated surface soils, depositing them in nearby floodplains.

Digging for road construction and repair also contributed to the drop by mixing and burying some contaminated soil deeper underground.

Foundation soils are less affected by these processes. Also, soils near many older homes continue to receive chips and dust from old leaded paint. Soil lead levels are thus decreasing more slowly near older homes, and those soils continue to pose a higher risk.

Traffic density also factors into the risk equation.

In larger cities that see more traffic, such as New York or Chicago, soil lead levels along heavily traveled streets have historically been high. In

larger cities, those levels may still be unacceptably high, Wade said. Implementing regular, widespread soil testing is the only way to know for certain.

"Large-scale sampling reveals patterns of soil lead distribution that you miss through spot checking," Richter said. "That can make the problem more predictable and remediable and encourage cities to act on this issue rather than just letting it be."

Wade and Richter's co-authors on the new study were Nancy Bao, Paul Heine, Mary Osteen and Kevin Tan, all of Duke, and Christopher Craft of Indiana University.

The study stemmed from a faculty-student research project that Wade and Richter led for Duke's Bass Connections program in 2018-2020.

More information: Anna M. Wade et al, Urban-Soil Pedogenesis Drives Contrasting Legacies of Lead from Paint and Gasoline in City Soil, *Environmental Science & Technology* (2021). [DOI: 10.1021/acs.est.1c00546](https://doi.org/10.1021/acs.est.1c00546)

Provided by Duke University School of Nursing

Citation: Lead levels in urban soil are declining but hotspots persist (2021, May 27) retrieved 12 May 2024 from <https://phys.org/news/2021-05-urban-soil-declining-hotspots-persist.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.