


Victoria's watch catchments may not recover from drought, study finds

May 13 2021



Percentage of the 161 study water catchments that displayed low runoff behaviour before, during and after the Millennium Drought from ~1997-2010. The shaded area from 2010 shows the percentage of water catchments that had not recovered from the drought. Credit: Monash University

Australian-first research by Monash University discredits the theory that rivers and underground water supplies eventually replenish following droughts or floods. Following the Australian Millennium Drought, one-third of Victoria's water catchments still had not recovered from drought nearly eight years later.

For those water catchments not recovered, roughly 80 per cent showed no evidence of recovering anytime soon.

One-third of the water catchments included in a Victorian study had not recovered from a [severe drought](#) nearly eight years later, Australian-first research from Monash University shows.

Globally, science holds the common view that rivers and underground water supplies eventually replenish following periods of severe [drought](#) or flood.

This study, led by Dr. Tim Peterson from Monash University's Department of Civil Engineering and published today in the prestigious international journal *Science*, is the first in the world to challenge this widely held view.

Researchers used statistical models of rainfall and streamflow at 161 water catchments across Victoria, each with over 30 years of data and no upstream dams or water extractions. The area surveyed is about the size of the United Kingdom or half that of the US state of California.

Dr. Peterson and research colleagues from The University of Melbourne discovered that when the drought ends, some rivers continue to behave like they're still in a drought for years afterwards and many have not yet recovered.

Specifically, the runoff, as a fraction of precipitation, had not recovered in 37 percent of water catchments in Victoria after Australia's Millennium Drought, and the number of recovering water catchments remained stagnant.

This means that 100mm of precipitation before the drought in 1990 created more river flow than the same 100mm in 2017, therefore delivering a 30 percent reduced streamflow after the drought.

The number of water catchments with a low or very low runoff state

increased rapidly from 1996 to the end of the meteorological drought in summer 2010. By 2011, only 15 percent of water catchments had recovered.

The Millennium Drought, regarded as one of the worst droughts to hit Australia in its modern history, crippled the Murray-Darling Basin and placed extreme pressure on ecosystems, agricultural production and urban water supply in the south-eastern part of the country. It ended with a La Nina weather event in 2010.

A water [catchment](#), or watershed, is any area of land that captures precipitation, which then flows into common outlets, such as a river, stream, bay or lake. Almost all of Victoria's water supply comes from streamflow.

Dr. Peterson said the regeneration of water catchments after severe drought had major implications for global long-term water resource planning and aquatic environments, especially when climate change is added on top of their findings.

"Our findings suggest hydrological droughts can persist indefinitely after meteorological droughts and that the mechanism for recovery remains an open question," Dr. Peterson said.

"This new discovery just appears to be the way catchments naturally behave. It's not explained by factors like land use. They are just more complex than we thought."

Each water catchment analysed for this study had at least 15, seven and five years of streamflow observations before, during and after the Millennium Drought respectively, and had no major upstream reservoirs or river extractions.

Across all 161 water catchments, researchers found eight years into the drought, 51 percent of the catchments switched into a low or very low runoff state. When the drought ended in 2010, primarily the eastern water catchments returned to a normal runoff state (see figure).

Importantly, by mid-2017, nearly eight years after the drought, more than one-third of water catchments still remained within a low runoff state, and have not recovered back to the pre-drought behaviour.

Dr. Peterson said evidence also suggested vegetation responded to the drought by increasing the fraction of precipitation going to transpiration—the process of water movement through a plant and its evaporation from leaves.

"Practically, this implies that in response to the Millennium Drought, vegetation in selected water catchments responded by maintaining similar rates of transpiration," he said.

Researchers say they've shown that water catchments are more complex than previously thought and that the findings are helping water agencies to better plan for the future.

Dr. Peterson and his co-authors at The University of Melbourne have been working with, and communicating the findings to, the Victorian and national water agencies; most recently through the broader findings of the Victorian Water and Climate Initiative.

He says: "it's exciting that the findings have already begun to be used in how water is managed. We are now developing mathematical tools to further help water management use these findings to ensure long-term [water](#) supply within a challenging and changing climate."

More information: T.J. Peterson at Monash University in Clayton,

VIC, Australia et al., "Watersheds may not recover from drought," *Science* (2021). [science.sciencemag.org/cgi/doi ... 1126/science.abd5085](https://science.sciencemag.org/cgi/doi/10.1126/science.abd5085)

"River basins on the edge of change," *Science* (2021).
[science.sciencemag.org/cgi/doi ... 1126/science.abi8770](https://science.sciencemag.org/cgi/doi/10.1126/science.abi8770)

Provided by Monash University

Citation: Victoria's watch catchments may not recover from drought, study finds (2021, May 13)
retrieved 3 May 2024 from
<https://phys.org/news/2021-05-victoria-catchments-recover-drought.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.