

Animals' ability to adapt their habitats key to survival amid climate change

June 18 2021



Michael Dillon (left), an associate professor in the University of Wyoming Department of Zoology and Physiology, and Arthur Woods (right), a professor of biological sciences at the University of Montana, were part of a research group that examined animals' ability to respond to climate change likely depends on how well they modify their habitats, such as nests and burrows. Here, Dillon bends over to examine a plant and measure microclimates at the UW-National



Park Service Research Station in Grand Teton National Park. Credit: Sylvain Pincebourde

Birds build nests to keep eggs and baby nestlings warm during cool weather, but also make adjustments in nest insulation in such a way the little ones can keep cool in very hot conditions. Mammals, such as rabbits or groundhogs, sleep or hibernate in underground burrows that provide stable, moderate temperatures and avoid above-ground conditions that often are far more extreme outside the burrow.

Michael Dillon, an associate professor in the University of Wyoming Department of Zoology and Physiology, was part of a research group that examined animals' ability to respond to <u>climate change</u> likely depends on how well they modify their habitats, such as nests and burrows.

So, how are these animals doing? Are they succeeding, struggling, or are their efforts a mixed bag in adapting their habitats to <u>climate</u> change?

"One of the key reasons that we wrote this paper is that we don't know the answer to this very important question!," Dillon says. "We hope the paper will encourage scientists to begin answering this question."

Dillon is a co-author of a paper, titled "Extended Phenotypes: Buffers or Amplifiers of Climate Change?," that was published June 16 in *Trends in Ecology & Evolution*. The journal publishes commissioned, peer-reviewed articles in all areas of ecology and evolutionary science.

The lead author of the paper is Arthur Woods, a professor of biological sciences at the University of Montana. Other contributors to the paper were from the University of Tours in Tours, France; and Stellenbosch



University in Stellenbosch, South Africa.

The study investigated extended phenotypes, which are modifications that organisms—birds, insects and mammals—make to their habitats.

"An extended phenotype can range from simply a hole in the ground occupied by an animal to leaves rolled into cavities by insects, to nests of all shapes and sizes built by birds and mammals, to <u>termite mounds</u> and bee colonies," Dillon says.

Extended phenotypes are important because they filter climate into local sets of conditions immediately around the organism. This is what biologists call the microclimate.

Because extended phenotypes are constructed structures, they often are modified in response to local climate variation and, potentially, in response to climate change. This process is called plasticity of the extended phenotype.

"One example might be a bird nest that is well insulated to protect eggs or young birds from cold. As climates warm, if the bird does not adjust insulation in the nest, it may, in fact, cause the young to overheat," Dillon explains.

In another prime example, termites build mounds that capture wind and solar energy to drive airflow through the colony, which stabilizes temperature, relative humidity and oxygen levels experienced by the colony.

However, the idea of microclimates is broader than constructed habitats. Microclimates typically differ substantially from nearby climates, which means that the climate in an area may provide little information about what animals experience in their microhabitats.



As an analogy, although a weather station might tell the public that the temperature in Laramie is 90 degrees Fahrenheit, simply by moving from the south to the north side of a building, one can experience microclimates that are strikingly different and often not captured by the weather data, Dillon says.

The same is true of animals of many different sizes. For example, a moose can move from an open sagebrush landscape to a shaded river corridor to cool off; a snake can move from its underground hole to a sunny rock to warm up; and a tiny insect shuttling between the top and bottom of a leaf can experience temperature differences of more than 20 degrees Fahrenheit.

"So, animals use microclimates, both by simply moving but also by building structures, such as nests, burrows, mounds and mines," Dillon says.

Across the globe, rising levels of carbon dioxide in the Earth's atmosphere are causing temperatures to rise and precipitation patterns to shift. For biologists, a key problem is to understand current effects of climate change on species, and to predict future effects, including how species' ranges may shift and what the relative risks of extinction are for different animal species' groups.

The research team favors a renewed effort to understand how extended phenotypes mediate how organisms experience climate change.

"We need a much better understanding of the basic biophysical principles by which extended phenotypes alter local conditions," says Sylvain Pincebourde, an ecologist in the Insect Biology Research Institute at the University of Tours and one of the paper's co-authors.

Another key challenge is to understand how much plasticity there is in



extended phenotypes, and how much and how rapidly they can evolve.

"At this point, we pretty much have no idea," Dillon says. "Can structures that buffer temperature variability keep up with the pace of climate change?"

More information: H. Arthur Woods et al, Extended phenotypes: buffers or amplifiers of climate change?, *Trends in Ecology & Evolution* (2021). DOI: 10.1016/j.tree.2021.05.010

Provided by University of Wyoming

Citation: Animals' ability to adapt their habitats key to survival amid climate change (2021, June 18) retrieved 9 April 2024 from https://phys.org/news/2021-06-animals-ability-habitats-key-survival.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.