


Bacteria hijack latent phage of competitor

June 10 2021



As virus particles, phages infect bacteria in order to ensure their own progress.
Credit: Thomas Böttcher

This targeted control of phages provides entirely new biotechnological and therapeutic approaches, e.g. for phage therapies. The results

produced in the context of an ERC grant have been published in the *Journal of the American Chemical Society*.

The human body and its microbiota harbor a large amount of phages. These infect bacteria as virus particles to ensure their own survival. One of their strategies is to integrate into the bacterial genome and multiply via bacterial cell division. However, external signal molecules can trigger the phages' sudden awakening from their dormant stage. Once activated, they destroy their host, the bacterium, and thus release their newly produced viral particles. With a prestigious ERC Consolidator Grant of the European Research Council, Thomas Böttcher investigates the switch from the sleeping (lysogenic) to the activated (lytic) lifestyle of phages.

Warfare between microbes

"We already know that phages decisively influence the population dynamics of bacteria and that microorganisms compete by using [chemical weapons](#)," says Thomas Böttcher, Professor of Microbial Biochemistry of the Faculty of Chemistry and the Centre of Microbiology and Environmental Systems Science. "We now wanted to investigate whether, in the complex microbial ecosystems, there are also microbes that specifically activate phages in order to use them against their competitors."

Indeed, the researchers could show that the bacterium *Pseudomonas aeruginosa* produces large amounts of a signal molecule that triggers the conversion of a [phage](#), residing in a strain of the species *Staphylococcus aureus*, from a quiet companion into a deadly parasite.

Highly selective phage activation

"We were completely surprised to find that the chemical compound

pyocyanin, which we were able to isolate and synthesize, only specifically activated one of several phages of *Staphylococcus aureus*. Pyocyanin is therefore a highly selective agent," says co-author Magdalena Jancheva.

The drug mitomycin C induces DNA damage in bacterial cells and causes phages to leave their dying host, but according to Thomas Böttcher, "it activates all phages in the bacteria in a non-selective manner." The researchers also observed that pyocyanin releases even more phages in *Staphylococcus aureus* than mitomycin C, pyocyanin therefore had an "remarkably strong effect."

Discovery provides new perspectives

The bacterial species *Pseudomonas aeruginosa* and *Staphylococcus aureus* occupy the same ecological niche in the [human body](#). As pathogens, they frequently occur in the lungs of patients with cystic fibrosis, a congenital metabolic disease. *Staphylococcus* bacteria dominate at a young age, while *Pseudomonas* bacteria become more prevalent with increasing age.

The current study demonstrates the efficiency of activation of latent phages by chemical signaling agents in the battle for space and resources between bacterial strains. It provides the first evidence that chemical signaling agents can exhibit selectivity for specific phages in a polylysogenic bacterial strain. Here, the activated phage (phiMBL3) revealed a previously unknown molecular switch through which the signaling agent acts.

"Certain [signaling molecules](#) could make it possible to combat pathogens via phage activation—they could thus be used to initiate internal phage therapy," Thomas Böttcher states. At the same time, the phages' molecular switches, which selectively trigger the production of viral particles through a signaling molecule such as pyocyanin, could also

serve as a new tool for biotechnology or synthetic biology. "Our findings open up a wide field in which we want to move forward," the researchers conclude.

More information: Magdalena Jancheva et al, A Metabolite of Pseudomonas Triggers Prophage-Selective Lysogenic to Lytic Conversion in Staphylococcus aureus, *Journal of the American Chemical Society* (2021). [DOI: 10.1021/jacs.1c01275](https://doi.org/10.1021/jacs.1c01275)

Provided by University of Vienna

Citation: Bacteria hijack latent phage of competitor (2021, June 10) retrieved 20 April 2024 from <https://phys.org/news/2021-06-bacteria-hijack-latent-phage-competitor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.