

## A 50% rise in carbon dioxide level could reduce rainfall in the Amazon more than deforestation

June 29 2021, by André Julião



Simulations run by Brazilian scientists on a supercomputer at the National Space Research Institute (INPE) show that the direct impact of rising levels of carbon dioxide over the Amazon rainforest would be a reduction in rainfall equivalent to or even greater than the impact of complete substitution of the forest by pasture. The result calls attention to the need for regional and global action to mitigate the adverse effects of climate change. Credit: João Marcos Rosa/AmazonFACE

A 50% rise in the level of carbon dioxide ( $CO_2$ ) in the atmosphere could reduce rainfall in the Amazon as much as or even more than substitution of the entire forest by pasture. The rise in  $CO_2$  would reduce the amount



of water vapor emitted by the forest, leading to a 12% annual drop in the volume of rainfall, while total deforestation would reduce rainfall by 9%.

These estimates are presented in a study published in *Biogeosciences* by scientists affiliated with the National Space Research Institute (INPE), the University of São Paulo (USP) and the University of Campinas (UNICAMP) in Brazil, and with Munich Technical University (TUM) in Germany.

"CO<sub>2</sub> is a basic input for photosynthesis, so when it increases in the atmosphere, plant physiology is affected and this can have a <u>cascade</u> <u>effect</u> on the transfer of moisture from trees to the atmosphere [transpiration], the formation of rain in the region, <u>forest</u> biomass, and several other processes," said David Montenegro Lapola, last author of the article.

Lapola is a professor at UNICAMP's Center for Meteorological and Climate Research Applied to Agriculture (CEPAGRI) and principal investigator for a project funded via the FAPESP Research Program on Global Climate Change (RPGCC). The study was also part of a Thematic Project funded by FAPESP and supported by a postdoctoral fellowship awarded to the penultimate author.

The researchers set out to investigate how the physiological effects of rising atmospheric  $CO_2$  on plants influence the rainfall regime. Plants transpire less as the supply of  $CO_2$  increases, emitting less moisture into the atmosphere and hence generating less rain.

Normally, however, predictions regarding the increase in atmospheric  $CO_2$  do not dissociate its physiological effects from its effects on the balance of radiation in the atmosphere. In the latter case, the gas prevents some of the Sun's reflected energy escaping from the



atmosphere, causing the warming phenomenon known as the greenhouse effect.

Projections presented in the latest report of the Intergovernmental Panel on Climate Change (IPCC), taking into account changes in the atmospheric radiation balance plus the physiological effects on plants, had already forecast a possible reduction of up to 20% in annual rainfall in the Amazon and shown that much of the change in the region's precipitation regime will be controlled by the way the forest responds physiologically to the increase in  $CO_2$ .

For the recently published study, the researchers ran simulations on the supercomputer at INPE's Center for Weather Forecasting and Climate Studies (CPTEC) in Cachoeira Paulista, state of São Paulo. They projected scenarios in which the atmospheric level of  $CO_2$  rose 50% and the forest was entirely replaced by pasture to find out how these changes affected the physiology of the forest over a 100-year period.

"To our surprise, just the physiological effect on the leaves of the forest would generate an annual fall of 12% in the amount of rain [252 millimeters less per year], whereas total deforestation would lead to a fall of 9% [183 mm]. These numbers are far higher than the natural variation in precipitation between one year and the next, which is 5%," Lapola said.

The findings draw attention to the need for local action to reduce deforestation in the nine countries that share the Amazon basin and global action to reduce  $CO_2$  emissions into the atmosphere by factories, vehicles and power plants, for example.

Lapola is one of the coordinators of the AmazonFACE experiment. The acronym stands for Free-Air Carbon dioxide Enrichment. Installed not far north of Manaus, the experiment will raise the level of  $CO_2$  over



small tracts of rainforest and analyze the resulting changes to <u>plant</u> <u>physiology</u> and the atmosphere. The experiment could anticipate the <u>climate change scenario predicted for this century.</u>

## **Transpiration in forest and pasture**

The scenarios projected by the computer simulations showed the decrease in rainfall being caused by a reduction of about 20% in leaf transpiration. The reasons for the reduction are different in each situation, however.

Stomata are microscopic portals in plant leaves that control gas exchange for photosynthesis. They open to capture  $CO_2$  and at the same time emit water vapor. In the scenario with more  $CO_2$  in the air, stomata remain open for less time and emit less water vapor, reducing cloud formation and rainfall.

Total leaf area shrinkage is another reason. If the entire forest were replaced by pasture, leaf area would shrink 66%. This is because the forest contains several layers of superimposed leaves in trees, so that leaf area per square meter is up to six times what it is on the ground. Lastly, both rising levels of  $CO_2$  and deforestation also influence the wind and the movement of air masses, which play a key role in the precipitation regime.

"The forest canopy has a complex surface made up of the tops of tall trees, low trees, leaves and branches. This is called canopy surface roughness. The wind produces turbulence, with eddies and vortices that in turn produce the instability that gives rise to the convection responsible for heavy equatorial rainfall," Lapola said. "Pasture has a smooth surface over which the wind always flows forward, and without forest doesn't produce vortices. The wind intensifies as a result, bearing away most of the precipitation westward, while much of eastern and



central Amazonia, the Brazilian part, has less rain."

The decrease in transpiration caused by rising levels of  $CO_2$  leads to a temperature increase of up to two degrees because there are fewer water droplets to mitigate the heat. This factor triggers a cascade of phenomena that result in less rain owing to inhibition of so-called deep convection (very tall rain clouds heavy with water vapor).

"A next step would be to test other computational models and compare the results with our findings," Lapola said. "Another important initiative would consist of more experiments like FACE, as only these can supply data to verify and refine modeling simulations like the ones we performed."

**More information:** Gilvan Sampaio et al,  $CO_2$  physiological effect can cause rainfall decrease as strong as large-scale deforestation in the Amazon, *Biogeosciences* (2021). <u>DOI: 10.5194/bg-18-2511-2021</u>

## Provided by FAPESP

Citation: A 50% rise in carbon dioxide level could reduce rainfall in the Amazon more than deforestation (2021, June 29) retrieved 3 May 2024 from <u>https://phys.org/news/2021-06-carbon-dioxide-rainfall-amazon-deforestation.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.