

The complex organization of an ant colony

June 18 2021



Credit: Unsplash/CC0 Public Domain

The colonies of social insects are complex systems that are entirely self-organized. Scientists who looked at the demographic, genetic and morphological structure of ant colonies were able to show how this self-organization works in practice. The study was supported by the Swiss National Science Foundation (SNSF) and is the subject of a paper

published in *PLOS Biology*.

Colonies of clonal raider [ants](#) (*Ooceraea biroi*) were used as model organisms in this study. This Asian predatory ant is interesting for researchers because they can easily control the age, genetic and morphological makeup of colonies.

In their study, they have shown that the organization in a homogeneous [colony](#) is modified when different individuals are introduced.

"Individuals of different sizes increased the division of labor in the colony, while genetically [different individuals](#) reduced it," explains Yuko Ulrich, the study's first author and former researcher at the University of Lausanne. "In fact, each source of heterogeneity generates a distinct pattern of behavioral organization in the colony," she says.

Who will do the dishes?

The results have surprised scientists because they go against some current theories on social groups. "It is thought that individuals act according to their tolerance threshold to stimuli," says Yuko Ulrich. To illustrate this model, the scientist proposes a comparison with humans: "In a family, some individuals react much more quickly than others to a pile of dirty dishes. They will therefore find themselves doing the dishes more often: a division of labor is taking place."

To explain the results of their ant study, the researchers had to expand this theoretical model. It now takes into account not only the stimulus tolerance but also the efficiency of each individual in performing a task and the overall workload of the colony.

This model still needs to be tested, but it is already opening up possibilities, explains Yuko Ulrich. It could provide a better understanding of other complex [biological systems](#) in which large

numbers of heterogeneous individuals interact; this in turn could be used to generate more accurate predictions of collective outcomes.

Automatic tracking of individuals

The scientists carried out their tests on 120 colonies of ants, homogeneous or heterogeneous, which they raised in transparent Petri dishes. In order to observe the behavior of each ant, they developed an automatic tracking system. "This is the first time such a system has been implemented on this scale in an ant study. Without this type of software, tracking would have been impossible," explains Yuko Ulrich. Each experiment lasted about a month. About 7000 images were taken for each colony.

In order to be identified by the software, each ant was marked with a unique color combination. Based on the spatial distribution of each individual, the software calculated an index of the division of labor in each colony. It does not specify what each individual does, but it gives clues to their role. "If an ant often stays near the nest, we can assume that it is caring for the larvae. An ant that moves around a lot is more likely to be in charge of foraging," explains the scientist.

More information: Yuko Ulrich et al, Response thresholds alone cannot explain empirical patterns of division of labor in social insects, *PLOS Biology* (2021). [DOI: 10.1371/journal.pbio.3001269](https://doi.org/10.1371/journal.pbio.3001269)

Provided by Swiss National Science Foundation

Citation: The complex organization of an ant colony (2021, June 18) retrieved 29 April 2024 from <https://phys.org/news/2021-06-complex-ant-colony.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.