

Global peatland dry-out could release 860 million tons of carbon dioxide per year

June 16 2021, by Yuanyuan Huang and Yingping Wang



Credit: Andrea Piacquadio from Pexels

Peatlands, such as fens, bogs, marshes and swamps, <u>cover just 3%</u> of the Earth's total land surface, yet store over one-third of the planet's soil carbon. That's more than the carbon stored in all other vegetation



combined, including the world's forests.

But peatlands worldwide are running short of water, and the amount of greenhouse gases this could set loose would be devastating for our efforts to curb <u>climate change</u>.

Specifically, <u>our new research</u> in Nature Climate Change found drying peatlands could release an additional 860 million tons of carbon <u>dioxide</u> into the atmosphere every year, by around 2100. To put this into perspective, Australia emitted <u>539 million tons in 2019</u>.

To stop this from happening, we need to urgently preserve and restore healthy, water-logged conditions in peatlands. These thirsty peatlands need water.

Peatlands are like natural archives

Peatlands are found across the world: the arctic tundra, coastal marshes, tropical swamp forests, mountainous fens and blanket bogs on subantarctic islands.

They're characterized by having water-logged soil filled with very slowly decaying <u>plant material</u> (the "peat") that accumulated over tens of thousands of years, preserved by the low-oxygen environment. This partially decomposed plant debris is locked up in the soils as organic carbon.





The giant dragonfly (Petalura gigantea) is listed as endangered under NSW environment law. Credit: <u>Christopher Brandis/iNaturalist</u>, <u>CC BY-NC</u>

Peatlands can act like natural archives, letting scientists and archaeologists reconstruct past climate, vegetation, and even <u>human lives</u>. In fact, an estimated <u>20,500 archaeological sites</u> are preserved under or within peat in the UK.

As unique habitats, peatlands are home for many native and endangered species of plants and animals that occur nowhere else, such as the whitebellied cinclodes (Cinclodes palliatus) in Peru and Australia's giant dragonfly (Petalura gigantea), the world's largest. They can also act as



migration corridors for birds and other animals, and can purify water, regulate floods, retain sediments and so on.

But over the past several decades, humans have been draining global peatlands for a range of uses. This includes planting trees and crops, harvesting peat to burn for heat, and for other land developments.

For example, some peatlands rely on groundwater, such as portions of <u>the Greater Everglades</u>, the largest freshwater marsh in the United States. Over-pumping groundwater for drinking or irrigation has cut off the peatlands' source of water.

Together with the regional drier climate due to global warming, our peatlands are drying out worldwide.

What happens when peatlands dry out?

When peat isn't covered by water, it could be exposed to enough oxygen to fuel aerobic microbes living within. The oxygen allows the microbes to grow extremely fast, enjoy the feast of carbon-rich food, and release carbon dioxide into the atmosphere.





A marsh in Les Sables d Olonne, France. Some peatlands are also a natural sources of methane, which is a more potent greenhouse gas than carbon dioxide. Credit: Arthur Gallois, Author provided

Some peatlands are also a natural source of methane, a potent greenhouse gas with the warming potential up to 100 times stronger than carbon dioxide.

But generating methane actually requires the opposite conditions to generating carbon dioxide. Methane is more frequently released in watersaturated conditions, while carbon dioxide emissions are mostly in unsaturated conditions.



This means if our peatlands are getting drier, we would have an increase in emissions of carbon dioxide, but a reduction in methane emissions.

So what's the net impact on our climate?

We were part of an international team of scientists across Australia, France, Germany, Netherlands, Switzerland, the US and China. Together, we collected and analyzed a large dataset from carefully designed and controlled experiments across 130 peatlands all over the world.

In these experiments, we reduced water under different climate, soil and environmental conditions and, using machine learning algorithms, disentangled the different responses of greenhouse gases.

Our results were striking. Across the peatlands we studied, we found reduced water greatly enhanced the loss of peat as carbon dioxide, with only a mild reduction of methane emissions.





A swamp forest in Peru. Credit: Rupesh Bhomia, Author provided

The net effect—carbon dioxide vs methane—would make our climate warmer. This will seriously hamper global efforts to keep temperature rise under 1.5°C.

This suggests if sustainable developments to restore these ecosystems aren't implemented in future, drying peatlands would add the equivalent of <u>860 million tons of carbon dioxide</u> to the atmosphere every year by 2100. This projection is for a "high emissions scenario", which assumes global greenhouse gas emissions aren't cut any further.



Protecting our peatlands

It's not too late to stop this from happening. In fact, many countries are already establishing <u>peatland</u> restoration projects.

For example, the <u>Central Kalimantan Peatlands Project</u> in Indonesia aims to rehabilitate these ecosystems by, for instance, damming drainage canals, revegetating areas with <u>native trees</u>, and improving local socioeconomic conditions and introducing more sustainable agricultural techniques.

Likewise, the Life Peat Restore project aims to restore 5,300 hectares of peatlands back to their natural function as <u>carbon</u> sinks across Poland, Germany and the Baltic states, over five years.

But protecting peatlands is a global issue. To effectively take care of our peatlands and our climate, we must work together urgently and efficiently.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Global peatland dry-out could release 860 million tons of carbon dioxide per year (2021, June 16) retrieved 27 June 2024 from <u>https://phys.org/news/2021-06-global-peatland-dry-out-million-tons.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.