

Imaging at the tip of a needle

June 18 2021




Fig. 1: Field transformations through multimode fibres. From: Memory effect

assisted imaging through multimode optical fibres

A team of physicists, led by Dr. David Phillips from the University of Exeter, have pioneered a new way in which to control light that has been scrambled by passage through a single hair-thin strand of optical fiber. These ultra-thin fibers hold much promise for the next generation of medical endoscopes—enabling high-resolution imaging deep inside the body at the tip of a needle.

Conventional endoscopes are millimeters wide and have limited resolution—so cannot be used to inspect individual cells. Single optical fibers are approximately 10x narrower and can enable much higher-resolution imaging—enough to examine the features of individual cells directly inside living tissue. It is normally only possible to view cells once they have been taken outside the body and placed in a microscope.

The catch is that we can't directly look through optical fibers, as they scramble the [light](#) sent through them. This problem can be solved by first calibrating an optical fiber to understand how it blurs images, and then using this calibration information as a key to decipher [images](#) from the scrambled light. Earlier this year, Dr. Phillips' group developed a way to measure this key extremely rapidly, in collaboration with researchers from Boston University in the U.S., and the Leibniz Institute of Photonic Technologies in Germany [paper: Compressively sampling the optical transmission matrix of a multimode fiber, published in *Light: Science and Applications*, April 21st 2021].




Fig. 2: The quasi-radial memory effect in MMFs. From: Memory effect assisted imaging through multimode optical fibres

However, the measured key is very fragile, and easily changes if the fiber bends or twists, rendering deployment of this technology in real clinical settings currently very challenging. To overcome this problem, the Exeter based team have now developed a new way to keep track of how the image unscrambling key changes while the fiber is in use. This provides a way to maintain high resolution imaging even as a single fiber based micro-endoscope flexes. The researchers achieved this by borrowing a concept used in astronomy to see through atmospheric turbulence and applying it to look through optical fibers. The method relies on a 'guide-star' – which in their case is a small brightly

fluorescing particle on the end of the fiber. Light from the guide-star encodes how the key changes when the fiber bends, thus ensuring imaging is not disrupted.

This is a key advance for the development of flexible ultra-thin endoscopes. Such imaging devices could be used to guide biopsy needles to the right place, and help identify diseased cells within the body.

Dr. Phillips, an Associate Professor in the Physics and Astronomy department at the University of Exeter, said: "We hope that our work brings the visualization of sub-cellular processes deep inside the body a step closer to reality—and helps to translate this technology from the lab to the clinic."

More information: Shuhui Li et al, Memory effect assisted imaging through multimode optical fibres, *Nature Communications* (2021). [DOI: 10.1038/s41467-021-23729-1](https://doi.org/10.1038/s41467-021-23729-1)

Provided by University of Exeter

Citation: Imaging at the tip of a needle (2021, June 18) retrieved 22 May 2024 from <https://phys.org/news/2021-06-imaging-needle.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--