

# Localized the gene for blue plum skin

June 9 2021

---



Japanese plums display a wide variety of skin and flesh colours both in hue and

pattern, going from anthocyanin-less green and yellow to anthocyanin-rich red, purple and blue. Credit: CRAG-IRTA

The presence and accumulation of the antioxidant pigment anthocyanin dictates fruit hue in plums, and the synthesis of this compound is known to be regulated by the MYB10 genes. Now, researchers from the Center for Research in Agricultural Genomics (CRAG) and the Institute of Agrifood Research and Technology (IRTA) have found the gene that determines Japanese plum skin color. In a study published at the scientific journal *Frontiers in Plant Science*, the team reveals that the plum genome contains several copies of the MYB10 genes, and that DNA variations in one of these copies cause plums to have anthocyanins in the skin (showing a blue to red color) or not (presenting a yellow or green tone).

The Japanese plum, appreciated for its juiciness, is the most abundant on the market for direct fresh consumption, and Spain is one of its largest producers in the European Union, with plum trees mainly grown in the regions of Extremadura, Andalusia and Murcia. This new study provides a highly efficient tool for early selection of colored and non-colored fruits in Japanese plum breeding programs, an advancement aligned with the goals of the International Year of Fruits and Vegetables (IYFV), designated by the UN General Assembly, to increase efficiency in fruit food systems and to promote healthy nutrition through fruit consumption.

## **Anthocyanins: more than color**

The red to blue hue of flowers and fruits is due to anthocyanins, a group of antioxidant pigments that enhance flower pollination and provide plants protection from light damage and dehydration. Including these

healthy antioxidants in our diet has been linked to anti-carcinogenic and anti-inflammatory effects, and to the prevention of cardiovascular diseases, diabetes and obesity.

Apples, pears, peaches, apricots, [plums](#), cherries and strawberries, all of which belong to the Rosaceae family, are broadly considered a source of anthocyanins, contained in their skin and flesh. As fruit color has both an important impact in consumer choice and nutritional quality, it is not surprising that there is considerable interest in breeding these crops to obtain new varieties bearing fruits with diverse colors, hues and patterns while enhancing their health benefits.



Red to blue-skinned plums, with a high accumulation of anthocyanins. Credit: CRAG-IRTA

## **Focusing on the Japanese plum**

Within the Rosaceae crops, Japanese plum is among those with the highest fruit color variation both in hue and pattern, going from anthocyanin-less green and yellow to red, purple or blue. "Previous studies in Rosaceae species show that the synthesis and accumulation of anthocyanins is regulated by the MYB10 [genes](#). Accordingly, the analysis of these genes in several Japanese plum has been proven to be an excellent model to understand how fruit color is determined," indicates Arnau Fiol, Ph.D. student at CRAG and first author of the article.

"In this study, we examined the MYB10 genes in a panel of Japanese plum varieties and we found that they are highly variable. Surprisingly, we discovered that some cultivars have three copies of one of the MYB10 genes, which adds up to the already complex analysis of plum color variation," explains Maria José Aranzana, IRTA researcher at CRAG in charge of this work. "By studying how these gene variants are inherited, we were able to identify which gene variant combinations are associated with the anthocyanin (red to blue) and anthocyanin-less (green or yellow) fruit skin colorations," she adds.

## **Boosting the development of new varieties**

Imagine we want to develop a new blue plum variety with lots of anthocyanins to benefit from its nutraceutical properties. New fruit tree varieties are obtained in breeding programs by crossing individuals and examining the hundreds or thousands of descendants looking for those

that present the desired trait, say a bluer plum, and that also meet demanding product quality standards. Between 10 and 20 years may pass from the first cross until the registration of a new variety, as it takes a long period for trees to bear fruits (about three to four years in plum trees) and they must pass several exhaustive evaluations.

"In our work, we have identified the gene variants that cause anthocyanins to accumulate or not in Japanese plum skin. This means that, by just studying the seedlings' DNA, we can efficiently predict the skin color of the fruits they will bear as soon as the plants germinate, allowing us to promptly discard all the seedlings that will produce green fruits. Thanks to this early screening, we'd need a smaller growing area and fewer resources (hydric, nutritional, phytosanitary and human) to obtain the desired new anthocyanin-rich plum variety, with the consequent economic and environmental benefits," points out Aranzana.

The reliable molecular marker for early selection of colored and non-colored Japanese plums developed in this research can be effectively used in breeding programs. Anticipating at a seedling level the color of the fruit that plants will produce in 3-4 years significantly shortens and optimizes the breeding process. Since the mechanism underlying variation in [fruit](#) color is conserved within the Rosaceae family, this tool holds promising prospects for anthocyanin-determined color selection in other related species of agronomic interest.

**More information:** Arnau Fiol et al, Characterization of Japanese Plum (*Prunus salicina*) PsMYB10 Alleles Reveals Structural Variation and Polymorphisms Correlating With Fruit Skin Color, *Frontiers in Plant Science* (2021). [DOI: 10.3389/fpls.2021.655267](https://doi.org/10.3389/fpls.2021.655267)

Provided by Centre for Research in Agricultural Genomics (CRAG)

Citation: Localized the gene for blue plum skin (2021, June 9) retrieved 19 April 2024 from <https://phys.org/news/2021-06-localized-gene-blue-plum-skin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.