

# How long-known genes continue to surprise researchers

June 16 2021




Fig. 1 From: Splicing and editing of ionotropic glutamate receptors: a comprehensive analysis based on human RNA-Seq data

The human genome was sequenced around 20 years ago. Since then, the sequence information encoding our proteins is known—at least in principle. However, this information is not continuously stored in the individual genes, but is divided into smaller coding sections. These

coding sections, also known as exons, are assembled in a process called splicing. Depending on the gene, different exon combinations are possible, which is why they are referred to as different or alternative splicing combinations.

Almost all 20,000 [human genes](#) can be alternatively spliced. A particularly huge variety of different splice variants is found in the brain, which allows for creating a huge diversity and allows to adapt the proteins to specific requirements. "However, it is not easy to determine, which [protein](#) variants are actually present," says Andreas Reiner. "Sequencing of already-spliced messenger RNAs (mRNAs), so-called RNA-Seq data, which are now increasingly being obtained with high-throughput approaches, offers a way out." Robin Herbrechter and Andreas Reiner now used such data to obtain an overview of all ionotropic glutamate receptor splice variants.

## **New glutamate receptor variants detected**

Using bioinformatic methods, the researchers aligned billions of mRNA sequence snippets to the genome to reconstruct the frequency of individual splice events. This method also enabled them to detect new, previously unknown splice variants. There were quite a few surprises: the [systematic analysis](#) showed that some variants found in the previously studied model organisms mouse and rat do not occur in humans at all, or are much less abundant than previously assumed.

"Among the newly identified isoforms, some are particularly exciting, as they are quite different from the previously known variants and could thus have new functions," says Robin Herbrechter. This includes a protein domain formed by the gene of the GluA4 AMPA receptor, as well as the first description of a delta receptor 1 (GluD1) isoform. While the focus will now be on analyzing these variants, also further bioinformatic analyses are planned, for example to determine which [cell](#)

[types](#) produce the different splice variants.

**More information:** Robin Herbrechter et al, Splicing and editing of ionotropic glutamate receptors: a comprehensive analysis based on human RNA-Seq data, *Cellular and Molecular Life Sciences* (2021).

[DOI: 10.1007/s00018-021-03865-z](https://doi.org/10.1007/s00018-021-03865-z)

Provided by Ruhr-Universitaet-Bochum

Citation: How long-known genes continue to surprise researchers (2021, June 16) retrieved 20 May 2024 from <https://phys.org/news/2021-06-long-known-genes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--