

Researchers resolve magnetic structures of different topological semimetals

June 24 2021, by Zhang Nannan




Figure 1. (a) Crystal and magnetic structures of PrAlGe; (b) M(H) curves up to high magnetic field of 30T; (c) angular-dependent magnetization; (d) critical isothermal analysis under high magnetic field. Credit: ZHANG Lei

Topological semimetals are one of the major discoveries in condensed-matter physics in recent years. The magnetic Weyl semimetal, in which the Weyl nodes can be generated and modulated by magnetization, provides an ideal platform for the investigation of the magnetic field-tunable link between Weyl physics and magnetism. But due to the lack of appropriate or high quality specimens, most of the theoretically expected magnetic topological semimetals have not been experimentally confirmed. Therefore, exploration of new magnetic topological semimetals is of great importance.

Recently, researchers from the High Magnetic Field Laboratory of the Hefei Institutes of Physical Science (HFIPS), in collaboration with researchers from Huazhong University of Science and Technology and Anhui University, resolved magnetic structures of different topological semimetals with the help of Resistive Magnet of China's Steady High Magnetic Field Facility (SHMFF) of HFIPS.

The team performed an investigation on high-quality single crystals of PrAlGe and DySb. For PrAlGe, the intrinsic ferromagnetic ordering acts as a Zeeman coupling to split the spin-up and spin-down bands, but the whole band structure is still kept. The study of magnetism suggested that the magnetic interaction in PrAlGe is of a 2D Ising type, revealing a uniaxial magnetic interaction along the c axis. However, the ordering moments are tilted from the c axis, which causes antiferromagnetism in the ab plane.

As for DySb, a field-induced tricritical phenomenon is revealed. Based on the magnetization analysis, a detailed H-T [phase diagram](#) around the phase transition is constructed when the [magnetic field](#) is applied along [001] direction.

This phase diagram is indicative of delicate competition and balance between multiple magnetic interactions in these systems and lays a solid

foundation for future research in topological transition and criticality.




Figure 2. H-T phase diagram for DySb. Credit: ZHANG Lei

More information: Wei Liu et al, Field-induced tricritical phenomenon and multiple phases in DySb, *Physical Review B* (2020). [DOI: 10.1103/PhysRevB.102.174417](https://doi.org/10.1103/PhysRevB.102.174417)

Wei Liu et al, Critical behavior of the magnetic Weyl semimetal PrAlGe, *Physical Review B* (2021). [DOI: 10.1103/PhysRevB.103.214401](https://doi.org/10.1103/PhysRevB.103.214401)

Provided by Chinese Academy of Sciences

Citation: Researchers resolve magnetic structures of different topological semimetals (2021, June 24) retrieved 9 February 2023 from

<https://phys.org/news/2021-06-magnetic-topological-semimetals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.