

# Microbes feast on crushed rock in subglacial lakes beneath Antarctica

June 29 2021

---



The field location including the tents and labs set up at Subglacial Lake Whillans, Antarctica. Credit: John Priscu

Pioneering research has revealed the erosion of ancient sediments found deep beneath Antarctic ice could be a vital and previously unknown source of nutrients and energy for abundant microbial life.

The study, led by the University of Bristol and published today in

*Communications Earth & Environment*, sheds new light on the many compounds supporting various microbes which form part of a huge subglacial ecosystem.

Lead author Dr. Beatriz Gill Olivas, a Post-Doctoral Research Associate at the University of Bristol, said: "Although the study focused on samples obtained from a single subglacial [lake](#), the results could have much wider implications. Subglacial Lake Whillans is part of a large interconnected hydrological system, so [erosion](#) taking place upstream could represent a potential source of biologically important compounds to this and other lakes within the system that might harbour thriving communities of microbial life."

The international research team replicated erosion processes in ice sheets by crushing sediments extracted from Lake Whillans, a subglacial lake in Antarctica, spanning around 60 km<sup>2</sup>, some 800m below the ice surface, then wetting these sediments and keeping them at 0 degrees Celsius without oxygen, to approximate subglacial conditions.

Findings showed a single crushing event down to a depth of 10 cm, followed by a 41-day incubation period, of these same sediments has the potential to provide up to a quarter (24 percent) of the estimated methane required by methanotrophs, tiny microbes which depend on methane as their source of carbon and energy, present in these environments. Substantial concentrations of hydrogen and carbon dioxide were also produced during crushing and incubation. These gases could potentially be used by methane-generating microbes, called methanogens, to produce enough methane which would help explain its widely varying levels within Lake Whillans.

Another previously unidentified finding was the detection of measurable concentrations of ammonium in water after incubation with crushed sediments. This is of particular significance to Lake Whillans, where

there is an abundance of microbial taxa that have the potential to derive energy from ammonium oxidation, a process known as nitrification. The study demonstrated a single high-energy crushing event, followed by the same incubation period, has the potential to produce more than the annual ammonium demand (120 percent) within the lake.

"Our previous understanding suggests the structure and function of subglacial ecosystems depends on the presence of redox species and redox gradients within liquid water. This study shows the process of erosion could potentially split water on freshly abraded mineral surfaces and produce hydrogen (a reducing end member species) and hydrogen peroxide (a highly oxidizing compound), creating a redox gradient and providing a previously unrecognised source of energy to microbial ecosystems," Dr. Gill Olivas explained.

"Only two previous studies have looked at the potential influence of erosion on subglacial energy and nutrient sources, which involved crushing largely unweathered rock samples. This is the first study to use highly weathered, ancient marine sediments, yet concentrations of gases measured largely agreed with previous results."

Dr. Gill Olivas added: "Whilst these experiments do not reflect the true extent of erosion under glaciers, they do hint at the potential for the crushing of sediments to supply important sources of nutriment to subglacial ecosystems. Further study is now needed to identify the complete range of reactions resulting directly from erosion processes and how they may vary with differences in the underlying [sediment](#)."

**More information:** Beatriz Gill-Olivas et al, Subglacial erosion has the potential to sustain microbial processes in Subglacial Lake Whillans, Antarctica, *Communications Earth & Environment* (2021). [DOI: 10.1038/s43247-021-00202-x](https://doi.org/10.1038/s43247-021-00202-x)

Provided by University of Bristol

Citation: Microbes feast on crushed rock in subglacial lakes beneath Antarctica (2021, June 29)  
retrieved 23 April 2024 from  
<https://phys.org/news/2021-06-microbes-feast-subglacial-lakes-beneath.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.