

Measuring the elimination of plastic particles from the body in mice

June 16 2021




Figure 1. PET-CT images of mice at 6, 12, 24, and 48 h after oral administration of 20 nm polystyrene particles (top row) and 1 µm polystyrene particles (bottom row). Green border: MIP, blue border: coronal slice, orange border: sagittal slice, and yellow border: transverse slice. Scale is percentage of injected dose per gram of tissue (%ID/g). Credit: Outi Keinänen et al.

The accumulation of micro- and nanoplastics in the environment continues at an alarming rate. A radiolabelling technique developed at the University of Helsinki made it possible to monitor the movement and accumulation of plastics in the mouse body, as well as their elimination from it.

Postdoctoral Researcher Outi Keinänen from the University of Helsinki developed a method to radiolabel [plastic particles](#) in order to observe

their biodistribution on the basis of radioactivity with the help of positron emission tomography (PET). As a radiochemist, Keinänen has in her previous radiopharmaceutical studies utilized PET imaging combined with computed tomography (CT), which produces a very accurate image of the anatomical location of the radioactivity signal.

In the recently completed study, radiolabelled plastic particles were fed to mice, and their elimination from the body was followed with PET-CT scans. This was the first time that the movement and location of plastic particles in a living mammalian system were observed in real time.

The study utilized polystyrene particles of four different sizes: 20 nm, 220 nm, 1 μm and 6 μm . The journey of the radiolabelled plastic particles through the [gastrointestinal tract](#) was followed for two days (48 hours) through PET-CT scans.

The study, which was recently published in *Scientific Reports* journal, demonstrated that most of the particles had been eliminated from the mice naturally, through feces within two days. Not much translocation of plastic particles from the gastrointestinal tract to elsewhere in the body was seen, and the smallest particles were eliminated from the body at a faster rate than the larger ones.

In addition to PET imaging, the findings were verified by thoroughly measuring the radioactivity of the tissues and organs of the mice. The persistence of the radiolabel on the surface of the plastic particles was verified by collecting murine gastrointestinal tracts at several different time points after administering the particles. The gastrointestinal tracts of mice that were put down at different timepoints were cut open, ground and separated into several fractions based on size. The share of the non-attached radiolabel was very small compared to the radiolabel still attached to the plastic particles. This was proof that the monitored radiation signal described the passage of the plastic particles well.

First and foremost, the study surveyed the usefulness of PET imaging in the study of micro- and nanoplastics, demonstrating that PET imaging enables accurate and non-invasive observation of plastic particles in living animals. Consequently, PET imaging may well become an important element of investigations into the health effects of plastics on mammals.

"While only a single small dose of polystyrene particles was fed to the mice, people are exposed daily to a range of micro- and nanoplastics. Therefore, we cannot draw direct conclusions on the accumulation of plastics in mammals and their effects on the basis of this study alone," Keinänen notes.

"In addition to ingesting plastics, the air we breathe contains small particles of plastic. Further studies are in fact in the pipeline," Keinänen promises.

Next up, the researchers wish to investigate the long-term consequences of daily exposure to micro- and nanoplastics, as well as the accumulation of inhaled plastic particles in mice. In future projects, the aim is to use different plastic materials in addition to polystyrene, the type of [plastic](#) used in this study.

More information: Outi Keinänen et al, Harnessing PET to track micro- and nanoplastics in vivo, *Scientific Reports* (2021). [DOI: 10.1038/s41598-021-90929-6](https://doi.org/10.1038/s41598-021-90929-6)

Provided by University of Helsinki

Citation: Measuring the elimination of plastic particles from the body in mice (2021, June 16) retrieved 3 May 2024 from <https://phys.org/news/2021-06-plastic-particles-body-mice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.