

Scientists report efficient use of waste paper for cellulase production by *Trichoderma longiflorum*

June 16 2021, by Zhang Nannan




Figure 1. Structural characterization of waste paper. Left: SEM ($\times 1000$ magnification) and EDS analysis of waste paper (A, waste office paper; B, tissue paper; C, magazine paper; D, corrugated board); Right: XRD spectra of different waste paper. Credit: *Journal of Cleaner Production*

Researchers at the Institute of Modern Physics (IMP) of the Chinese Academy of Sciences (CAS) have recently found that waste paper could be efficiently used for enzyme production by *Trichoderma longiflorum*. The study was published in *Journal of Cleaner Production*.

Cellulase, accounting for the world's third-largest share of the industrial

[enzyme](#) market behind amylase and protease, is widely used in biofuel, food, papermaking, textile washing, medicine, [animal feed](#) and agricultural waste treatment. However, the high production cost of enzyme remains a major challenge for its further industrial applications.

Waste paper, a recyclable waste resource, has high potential for industry due to its richness in cellulose, low cost and high abundance. Thus, to produce cellulase and xylanase by using [waste paper](#) not only reduces the cost of [enzyme production](#), but also achieves the efficient use of waste paper resources.

In the experiment, the researchers used waste office paper, corrugated board, magazine paper and tissue paper for enzyme production by *T. longiflorum* LC-M4, a mutant generated by heavy ion irradiation.

By studying the structural characterization of waste paper, they revealed that the cellulose is the main component in different waste paper sources, and the fillers such as [calcium carbonate](#) can provide nutrient elements for microbial growth and enzyme synthesis.




Figure 2. Enzyme activities after 192h of fermentation by *T. longiflorum* LC-M4 using waste paper and wheat bran as a mixed carbon source. Error bars indicate standard deviation. Credit: *Journal of Cleaner Production*

They found that the highest filter paper activities were achieved by using waste office paper and corrugated board, respectively. Moreover, the maximum β -glucosidase and xylanase activities were obtained from corrugated board substrates.

In addition, to further improve the enzyme activities, the researchers used the corrugated board as a mixed carbon source with wheat bran. Higher β -glucosidase activities were obtained, which were significantly

increased by 56.86% when compared with the corrugated board as the single carbon source.

The results confirmed that the corrugated board and wheat bran have great application potential as carbon sources for enzyme production.

These findings obtained from the study revealed the feasibility of utilizing the waste paper as an inductive carbon source for enzyme production, which is of great benefit both for the economy and environment.

More information: Miaoyin Dong et al, Efficient utilization of waste paper as an inductive feedstock for simultaneous production of cellulase and xylanase by *Trichoderma longiflorum*, *Journal of Cleaner Production* (2021). [DOI: 10.1016/j.jclepro.2021.127287](https://doi.org/10.1016/j.jclepro.2021.127287)

Provided by Chinese Academy of Sciences

Citation: Scientists report efficient use of waste paper for cellulase production by *Trichoderma longiflorum* (2021, June 16) retrieved 15 June 2024 from <https://phys.org/news/2021-06-scientists-efficient-paper-cellulase-production.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--