

Space, exercise may be critical to drylot beef heifer reproduction

June 24 2021, by Kay Ledbetter



Heifers lounge in a drylot. Credit: Texas A&M AgriLife



Space and exercise could be almost as important as food and water to the successful development of beef heifers raised in drylots, and quantifying that importance is the aim of a planned study by a Texas A&M University Department of Animal Science researcher in the College of Agriculture and Life Sciences.

Reproductive development of replacement heifers dictates the overall efficiency of cow-calf operations. In order to maximize efficiency and minimize resource use, beef industry replacement heifers are increasingly being housed in drylots. While the practice is less common in Texas than in other areas of the country, land-use trends suggest it may become more prevalent, said Texas A&M's Reinaldo Cooke, Ph.D., beef cattle production associate professor.

"As an example, Houston, Navasota and College Station will soon become one big metropolitan area," Cooke said. "We are going to have less grass resources for cattle, competing with urban development and crop production. I don't believe the whole industry will change to confined operations, but we need to find management systems to make sure we are maintaining or promoting production efficiency and fostering <u>animal welfare</u>."

That is why, he said, management and stocking <u>density</u> guidelines for heifers reared in drylots are urgently needed. Optimal nutritional programs have been developed and disseminated, but not those for heifer welfare aspects, including stocking density. These stocking density guidelines exist for poultry, swine and dairy cows, yet there is a dearth of information regarding stocking density for beef cattle, Cooke said.

He will address this critical need with a Texas A&M Department of Animal Science team consisting of animal welfare specialist Courtney Daigle, Ph.D., and reproductive physiologists Ky Pohler, Ph.D., Rodolfo



Cardoso, Ph.D., and Cliff Lamb, Ph.D., who is also department chair.

The team received a \$500,000 grant from the U.S. Department of Agriculture, USDA, National Institute of Food and Agriculture to look at stocking density and determine if management of stocking density can prevent development of the reproductive pathways needed for the heifer to reach puberty.

Researching for answers

In typical U.S. spring-calving herds, replacement heifers are weaned in the fall and exposed to their first breeding season the following spring at about 15 months of age. During late fall and winter, heifers are often moved to drylot systems to ensure adequate feeding for growth, with special care to their nutritional regimen.

"Also, it's becoming more common in the western U.S., especially where the cows and calves are grazing forage on public lands, in October to wean the calves and bring them into a drylot system to feed for the winter. Even in operations with normal grazing, when the winters are severe, the heifers are moved to drylots to facilitate management and feeding. This typically begins during a critical period of growth around 7-9 months."

Cooke said his research group is the first to investigate and portray the potential adversities that result from this management scheme to heifer welfare and reproductive development.

The current stocking and spacing recommendations are mostly related to feedlot cattle, and aren't adaptable to cow-calf systems, he said. With more operations opting to rear heifers in drylots to meet environmental challenges and limited resources, producers need specific guidelines regarding stocking density.



Cooke said his earlier studies have found that beef heifers moved into drylots had lower reproductive efficiency than expected.

"Every time we have heifers enclosed in a confined environment, their reproduction efficiency is less than heifers in open pasture that had a lower quality diet," he said. "The drylot heifers gained more weight with better food, but not reproductive efficiency."

Cooke said while looking at the literature to understand what was happening, he found no one had studied the reproductive differences between the pasture-fed and drylot-reared heifers.

"That prompted us to run this project to determine why puberty is being blocked."

Theory: Heifers need room to move

Cooke said heifers in drylots don't have much opportunity to exercise, which is important to their hormonal welfare. So, their study design will help determine if providing areas for voluntary exercise and more loafing space mitigates the negative effects of drylot-housing.

"We know exercise areas are often implemented in confined dairy operations to allow cows to exercise voluntarily, with positive outcomes to their wellbeing and productivity," he said. "So, we believe that providing heifers the opportunity to exercise will alleviate the detrimental effects of a high stocking density on their welfare and reproductive responses."

Additionally, the Texas A&M team will compare different stocking densities or drylot dimensions. Their theory: increased drylot dimensions accommodate larger group sizes, which also promotes physical activity and mitigates social stressors.



"We will have exercising, 30 minutes a day, and non-exercising heifers to determine if their reproductive inefficiency is a fact of being confined or the lack of exercise. Our second and third objectives will look at stocking densities—how many animals in the pens."

The only density guidelines they can use are from feedlots, but even those are from the 1980s, Cooke said. The Federation of Animal Science Societies recommends about 15 square yards per heifer in unpaved drylots.

"We will begin with the minimum space they suggest and work up to see when we reach the same level of reproduction as the heifers on pasture," he said.

Stocking density is a management decision that substantially affects livestock, Cooke said. The results of this study will help develop guidelines to promote the welfare and productivity of beef heifers reared in drylot systems.

Provided by Texas A&M University

Citation: Space, exercise may be critical to drylot beef heifer reproduction (2021, June 24) retrieved 28 July 2024 from <u>https://phys.org/news/2021-06-space-critical-drylot-beef-heifer.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.