


The sun's clock: New calculations support and expand planetary hypothesis

June 14 2021



The sun sported about a dozen active regions over a five-day period in May 2015. The bright, spindly strands that extend out of these active regions are particles spinning along magnetic field lines that connect areas of opposite polarity. Credit: Solar Dynamics Observatory, NASA

Solar physicists around the world have long been searching for satisfactory explanations for the sun's many cyclical, overlapping activity fluctuations. In addition to the most famous, approximately 11-year "Schwabe cycle", the sun also exhibits longer fluctuations, ranging from hundreds to thousands of years. It follows, for example, the "Gleissberg cycle" (about 85 years), the "Suess-de Vries cycle" (about 200 years) and the quasi-cycle of "Bond events" (about 1500 years), each named after their discoverers. It is undisputed that the solar magnetic field controls these activity fluctuations.

Explanations and models in expert circles partly diverge widely as to why the magnetic field changes at all. Is the sun controlled externally or does the reason for the many cycles lie in special peculiarities of the solar dynamo itself? HZDR researcher Frank Stefani and his colleagues have been searching for answers for years—mainly to the very controversial question as to whether the planets play a role in solar activity.

Rosette-shaped movement of the sun can produce a 193-year cycle

The researchers have most recently taken a closer look at the sun's orbital movement. The sun does not remain fixed at the center of the solar system: It performs a kind of dance in the common gravitational field with the massive planets Jupiter and Saturn—at a rate of 19.86 years. We know from the Earth that spinning around in its orbit triggers

small motions in the Earth's liquid core. Something similar also occurs within the sun, but this has so far been neglected with regard to its magnetic field.

The researchers came up with the idea that part of the sun's angular orbital momentum could be transferred to its rotation and thus affect the internal dynamo process that produces the [solar magnetic field](#). Such coupling would be sufficient to change the extremely sensitive magnetic storage capacity of the tachocline, a transition region between different types of energy transport in the sun's interior. "The coiled magnetic fields could then more easily snap to the sun's surface," says Stefani.

The researchers integrated one such rhythmic perturbation of the tachocline into their previous model calculations of a typical solar dynamo, and they were thus able to reproduce several cyclical phenomena that were known from observations. What was most remarkable was that, in addition to the 11.07-year Schwabe [cycle](#) they had already modeled in previous work, the strength of the magnetic field now also changed at a rate of 193 years—this could be the sun's Suess-de Vries cycle, which from observations has been reported to be 180 to 230 years. Mathematically, the 193 years arise as what is known as a beat period between the 19.86-year cycle and the twofold Schwabe cycle, also called the Hale cycle. The Suess-de Vries cycle would thus be the result of a combination of two external "clocks": the planets' tidal forces and the sun's own movement in the solar system's gravitational field.

Planets as a metronome

For the 11.07-year cycle, Stefani and his researchers had previously found strong statistical evidence that it must follow an external clock. They linked this "clock" to the tidal forces of the planets Venus, Earth and Jupiter. Their effect is greatest when the planets are aligned: a constellation that occurs every 11.07 years. As for the 193-year cycle, a

sensitive physical effect was also decisive here in order to trigger a sufficient effect of the weak tidal forces of the planets on the solar dynamo.

After initial skepticism toward the planetary hypothesis, Stefani now assumes that these connections are not coincidental. "If the sun was playing a trick on us here, then it would be with incredible perfection. Or, in fact, we have a first inkling of a complete picture of the short and long solar activity cycles." In fact, the current results also retroactively reaffirm that the 11-year cycle must be a timed process. Otherwise, the occurrence of a beat period would be mathematically impossible.

Tippling into chaos: 1000-2000-year collapses are not more accurately predictable

In addition to the rather shorter activity cycles, the sun also exhibits long-term trends in the thousand-year range. These are characterized by prolonged drops in activity, known as "minima", such as the most recent "Maunder Minimum", which occurred between 1645 and 1715 during the "Little Ice Age". By statistically analyzing the observed minima, the researchers could show that these are not cyclical processes, but that their occurrence at intervals of approximately one to two thousand years follows a mathematical random process.

To verify this in a model, the researchers expanded their solar dynamo simulations to a longer period of 30,000 years. In fact, in addition to the shorter cycles, there were irregular, sudden drops in magnetic activity every 1000 to 2000 years. "We see in our simulations how a north-south asymmetry forms, which eventually becomes too strong and goes out of sync until everything collapses. The system tips into chaos and then takes a while to get back into sync again," says Stefani. But this result also means that very long-term solar activity forecasts—for example, to

determine influence on climate developments—are almost impossible.

More information: F. Stefani et al, Shaken and Stirred: When Bond Meets Suess–de Vries and Gnevyshev–Ohl, *Solar Physics* (2021). [DOI: 10.1007/s11207-021-01822-4](https://doi.org/10.1007/s11207-021-01822-4)

Provided by Helmholtz-Zentrum Dresden-Rossendorf

Citation: The sun's clock: New calculations support and expand planetary hypothesis (2021, June 14) retrieved 21 June 2024 from <https://phys.org/news/2021-06-sun-clock-planetary-hypothesis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.