


The discovery of a new type of supernova illuminates a medieval mystery

June 28 2021



Las Cumbres Observatory and Hubble Space Telescope color composite of the electron-capture supernova 2018zd (the large white dot on the right) and the host starburst galaxy NGC 2146 (toward the left). Credit: NASA/STSCI/J. Depasquale; Las Cumbres Observatory

A worldwide team led by UC Santa Barbara scientists at Las Cumbres

Observatory has discovered the first convincing evidence for a new type of stellar explosion—an electron-capture supernova. While they have been theorized for 40 years, real-world examples have been elusive. They are thought to arise from the explosions of massive super-asymptotic giant branch (SAGB) stars, for which there has also been scant evidence. The discovery, published in *Nature Astronomy*, also sheds new light on the thousand-year mystery of the supernova from A.D. 1054 that was visible all over the world in the daytime, before eventually becoming the Crab Nebula.


Historically, supernovae have fallen into two main types: thermonuclear and iron-core collapse. A thermonuclear supernova is the explosion of a white dwarf star after it gains matter in a binary star system. These white dwarfs are the dense cores of ash that remain after a low-mass star (one up to about 8 times the mass of the sun) reaches the end of its life. An iron core-collapse supernova occurs when a massive star—one more than about 10 times the mass of the sun—runs out of nuclear fuel and its iron core collapses, creating a black hole or neutron star. Between these two main types of supernovae are electron-capture supernovae. These stars stop fusion when their cores are made of oxygen, neon and magnesium; they aren't massive enough to create iron.

While gravity is always trying to crush a star, what keeps most stars from collapsing is either ongoing fusion or, in cores where fusion has stopped, the fact that you can't pack the atoms any tighter. In an electron capture supernova, some of the electrons in the oxygen-neon-magnesium core get smashed into their atomic nuclei in a process called electron capture. This removal of electrons causes the core of the star to buckle under its own weight and collapse, resulting in an electron-capture supernova.

If the star had been slightly heavier, the core elements could have fused to create heavier elements, prolonging its life. So it is a kind of reverse Goldilocks situation: The star isn't light enough to escape its core

collapsing, nor is it heavy enough to prolong its life and die later via different means.

That's the theory that was formulated beginning in 1980 by Ken'ichi Nomoto of the University of Tokyo and others. Over the decades, theorists have formulated predictions of what to look for in an electron-capture supernova and their SAGB star progenitors. The stars should have a lot of mass, lose much of it before exploding, and this mass near the dying star should be of an unusual chemical composition. Then the electron-capture supernova should be weak, have little radioactive fallout, and have neutron-rich elements in the core.



Artist impressions of a super-asymptotic giant branch star and its core made up

of oxygen, neon and magnesium. This is the end state of stars around 8-10 solar masses, whose core is pressure supported by electrons. When the core becomes dense enough, neon and magnesium start to eat up electrons, reducing the core pressure and inducing a core-collapse supernova explosion. Credit: S. Wilkinson; Las Cumbres Observatory

The new study is led by Daichi Hiramatsu, a graduate student at UC Santa Barbara and Las Cumbres Observatory (LCO). Hiramatsu is a core member of the Global Supernova Project, a worldwide team of scientists using dozens of telescopes around and above the globe. The team found that the supernova SN 2018zd had many unusual characteristics, some of which were seen for the first time in a supernova.

It helped that the supernova was relatively nearby—only 31 million light-years away—in the galaxy NGC 2146. This allowed the team to examine archival images taken by the Hubble Space Telescope prior to the explosion and to detect the likely progenitor star before it exploded. The observations were consistent with another recently identified SAGB star in the Milky Way, but inconsistent with models of red supergiants, the progenitors of normal iron core-collapse supernovae.

The authors looked through all published data on supernovae, and found that while some had a few of the indicators predicted for electron-capture supernovae, only SN 2018zd had all six: an apparent SAGB progenitor, strong pre-supernova mass loss, an unusual stellar chemical composition, a weak explosion, little radioactivity and a neutron-rich core.

"We started by asking 'what's this weirdo?'" Hiramatsu said. "Then we examined every aspect of SN 2018zd and realized that all of them can be explained in the electron-capture scenario."

The new discoveries also illuminate some mysteries of the most famous supernova of the past. In A.D. 1054 a supernova happened in the Milky Way Galaxy that, according to Chinese and Japanese records, was so bright that it could be seen in the daytime for 23 days, and at night for nearly two years. The resulting remnant, the Crab Nebula, has been studied in great detail.



This composite image of the Crab Nebula was assembled by combining data from five telescopes spanning nearly the entire breadth of the electromagnetic spectrum. Credit: NASA, ESA, NRAO/AUI/NSF and G. Dubner (University of Buenos Aires)

The Crab Nebula was previously the best candidate for an electron-capture supernova, but its status was uncertain partly because the explosion happened nearly a thousand years ago. The new result increases the confidence that the historic SN 1054 was an electron-capture supernova. It also explains why that supernova was relatively bright compared to the models: Its luminosity was probably artificially enhanced by the supernova ejecta colliding with material cast off by the progenitor star as was seen in SN 2018zd.

Ken Nomoto at the Kavli IPMU of the University of Tokyo expressed excitement that his theory had been confirmed. "I am very pleased that the electron-capture supernova was finally discovered, which my colleagues and I predicted to exist and have a connection to the Crab Nebula 40 years ago," he said. "I very much appreciate the great efforts involved in obtaining these observations. This is a wonderful case of the combination of observations and theory."

Hiramatsu added, "It was such a 'Eureka moment' for all of us that we can contribute to closing the 40-year-old theoretical loop, and for me personally because my career in astronomy started when I looked at the stunning pictures of the Universe in the high school library, one of which was the iconic Crab Nebula taken by the Hubble Space Telescope."

"The term Rosetta Stone is used too often as an analogy when we find a new astrophysical object," said Andrew Howell, a staff scientist at Las Cumbres Observatory and adjunct faculty at UCSB, "but in this case I think it is fitting. This supernova is literally helping us decode thousand-year-old records from cultures all over the world. And it is helping us associate one thing we don't fully understand, the Crab Nebula, with another thing we have incredible modern records of, this supernova. In the process it is teaching us about fundamental physics: how some neutron stars get made, how extreme stars live and die, and about how the elements we're made of get created and scattered around the

universe." Howell also is the leader of the Global Supernova Project, and lead author Hiramatsu 's Ph.D. advisor.

More information: The electron-capture origin of supernova 2018zd, *Nature Astronomy* (2021). DOI: [10.1038/s41550-021-01384-2](https://doi.org/10.1038/s41550-021-01384-2) , www.nature.com/articles/s41550-021-01384-2

Provided by University of California - Santa Barbara

Citation: The discovery of a new type of supernova illuminates a medieval mystery (2021, June 28) retrieved 13 August 2024 from <https://phys.org/news/2021-06-supernova.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.