

# Vegetation growth in Northern Hemisphere is stunted by water constraints in warming climate

June 24 2021

---



The degraded vegetation in the Horqin Sand Land in Inner Mongolia, China.  
Credit: Lixin Wang, IUPUI

A first-of-its-kind large-scale study of vegetation growth in the Northern Hemisphere over the past 30 years has found that vegetation is becoming increasingly water-limited as global temperatures increase.

The results are significant since vegetation is one of the biggest factors when it comes to controlling water and [carbon](#) cycling across Earth, which influences global temperatures. The work by IUPUI and Indiana University Bloomington researchers Wenzhe Jiao, Lixin Wang, Qing Chang and Honglang Wang was published in the journal *Nature Communications* on June 18.

"Without water, living things struggle to survive, including plants," said Lixin Wang, senior author of the study and an associate professor of earth sciences at the School of Science at IUPUI. His ecohydrology group led the study. "Changes in [vegetation response](#) to [water availability](#) can result in significant shifts of climate-carbon interaction."

Honglang Wang is an assistant professor of statistics at the School of Science at IUPUI. Wenzhe Jiao, the first author, and Qing Chang are Ph.D. students at IUPUI and IU Bloomington, respectively.

This multidisciplinary research between the School of Science at IUPUI, the O'Neill School of Public and Environmental Affairs at IU Bloomington and two other universities began three years ago to determine vegetation constraints on a global scale. Until now, it was largely unknown, despite the growing interest in predicting global and regional trends in vegetation growth in response to [climate change](#).



The saguaro cactus (*Carnegiea gigantea*) of the Sonoran Desert in the Southwest United States. Credit: Lixin Wang, IUPUI


"Global temperature and the concentration of atmospheric CO<sub>2</sub>, or carbon dioxide, have been increasing," Lixin Wang said. These changes are expected to cause increased atmospheric water demand, more frequent extreme hot days, and drought events. All these factors indicate that vegetation growth may have suffered more and more water stress under a warming climate.

"However, quantifying the changes in vegetation constraints at large spatial and temporal scales is challenging," he said.

To overcome this obstacle, the researchers used satellite remote sensing data and meteorology data covering large spatial scales from 1982 to 2015.

"We developed our own metrics to indicate water constraints and then examined the changes in the metrics," Jiao said. "The study is quite computationally extensive since we examined the relationship between vegetation growth and water deficit at each grid cell over the whole extratropical Northern Hemisphere—604,800 data points each year—over more than 30 years."

The [data analysis](#) provided strong evidence of a widespread, significant increase in water vegetation constraint in the Northern Hemisphere over the studied period. Some regions, like the Great Plains in the United States, were comparatively worse than others.



For the entire study period, the vertical axis of the color legend is the trend of correlation coefficient for the 30 five-year moving windows. No color indicates unvegetated regions. Chartreuse stands for vegetation water surplus regions where surplus has been decreasing; navy indicates regions that have been experiencing an increase in water surplus; magenta is for water deficit regions that have been seeing an increase in water deficit; and regions colored yellow are

characterized by water deficit and a decrease in water deficit. Credit: Lixin Wang et. al., IUPUI

Until recently, elevated carbon in the atmosphere had increased plant growth, which has the benefit of removing more carbon from the atmosphere. However, this study reveals a cause for concern.

"Increasing water constraints on vegetation productivity may drive a shift from a period of increasing land carbon sink strength to a period in which climate change is reducing land carbon sink strength," Lixin Wang said.

In other words, the warming climate is increasing water constraints, reversing the earlier trend of stronger vegetation carbon uptake.

"Our research shows that increasing water constraints will likely limit continuous vegetation growth, thus slowing down the removal of CO<sub>2</sub> from the atmosphere by plants," Jiao said.

"The results emphasize the need for actions that could slow down CO<sub>2</sub> emissions," Lixin Wang said. "Without that, [water](#) constraints impacting plant growth—and the weakening of vegetation's ability to removal of CO<sub>2</sub> from the atmosphere—are unlikely to slow."

**More information:** Wenzhe Jiao et al, Observed increasing water constraint on vegetation growth over the last three decades, *Nature Communications* (2021). [DOI: 10.1038/s41467-021-24016-9](https://doi.org/10.1038/s41467-021-24016-9)

Provided by Indiana University

Citation: Vegetation growth in Northern Hemisphere is stunted by water constraints in warming climate (2021, June 24) retrieved 2 May 2024 from <https://phys.org/news/2021-06-vegetation-growth-northern-hemisphere-stunted.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.