

## Arginine methylation in global control of RNA splicing and translation

July 19 2021, by Liu Jia







Fig. 1. Systematic identification of PRMT interactome. (a) Schematic diagram showing PRMT1 to PRMT9. The light blue boxes represent the catalytic domains, the cyan and yellow boxes represent double E motifs and THW loop motifs that are specific to PRMTs, respectively. (b) Workflow for identifying PRMT-interacting proteins using BioID. (c) The biotinylated PRMT-interacting proteins detected by Western blotting using streptavidin-HRP (left) and silver staining (right). (d) The Venn diagram illustrating the PRMT interactome obtained from this study compared with methylarginine-containing proteins identified in the study by Larsen et al. (2016) and Guo et al. (2014). The correlation is calculated by Fisher's exact test, with the whole genome as the background. (e) Overlap of the interactome among different PRMTs. Fishers' exact test is used to calculate the P-value of the overlap. The PRMT interactome sare clustered by overlap significance (shown in red), and the numbers of overlapped protein are indicated in blue. Credit: *Science Bulletin* (2021). DOI: 10.1016/j.scib.2021.01.004

Thousands of proteins undergo arginine N-methylation, a widespread post-translational modification catalyzed by several protein arginine methyltransferases (PRMTs) that covalently link methyl groups to arginine side chains. Nine PRMTs, PRMT1 to PRMT9, have been identified in the human genome, which are further classified into three types according to the final methylarginine products. Arginine methylation plays key roles in various cellular processes.

However, global understanding of their molecular mechanisms and biological functions is limited due to the lack of a complete picture of the catalytic network for each PRMT as well as the absence of reliable antibodies and mechanically well-defined "erasers" and "readers".

In a study published in *Science Bulletin*, the researchers from the Shanghai Institute of Nutrition and Health (SINH) of the Chinese Academy of Sciences (CAS) identified the interacting proteins for all



human PRMTs, and demonstrated their functional importance in mRNA splicing and translation.

Using a sensitive BioID technology, the researchers systematically characterized the interactome and the <u>substrate specificity</u> of all known human PRMTs. They demonstrated significant overlapping of interactomes of human PRMTs with the known methylargininecontaining proteins. Different PRMTs are functionally redundant with a high degree of overlap in their substrates and high similarities between their putative methylation motifs.

Through <u>bioinformatics analysis</u> and experimental validation, the researchers revealed that RNA-binding proteins (RBPs) involved in regulating RNA splicing and translation are highly enriched in PRMT interactomes and undergo extensive arginine methylation, indicating their importance in regulating RNA metabolism.

By applying RNA-seq and Ribo-seq, they demonstrated that the inhibition of PRMTs leads to global alteration of splicing and translation, providing a genome-wide picture of the general functions of protein arginine methylation. Interestingly, they found that the inhibition of PRMT activity did not change the distribution of ribosome occupancy in different regions of mRNAs, suggesting that PRMTs affected the maturation of ribosomes before they are assembled onto the mRNAs.

Using RPS2 as an example, the researchers found that mutations in RPS2's methylation sites suppress ribosome assembly, translation, and eventually <u>cell growth</u>, indicating that arginine methylation of ribosomal proteins is critical to ribosomal assembly. This might be the main mechanistic reason for why PRMT affect translation.

This study provides new insights into <u>biological functions</u> of PRMTs and links individual PRMTs to their arginine methylation events, revealing



critical functions of arginine methylation in regulating RNA splicing and translation.

**More information:** Huan-Huan Wei et al, A systematic survey of PRMT interactomes reveals the key roles of arginine methylation in the global control of RNA splicing and translation, *Science Bulletin* (2021). DOI: 10.1016/j.scib.2021.01.004

Provided by Chinese Academy of Sciences

Citation: Arginine methylation in global control of RNA splicing and translation (2021, July 19) retrieved 25 June 2024 from <u>https://phys.org/news/2021-07-arginine-methylation-global-rna-splicing.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.