


# Synthesizing aryl sulfides from non-smelling, non-toxic compounds

July 21 2021



Ni-Catalyzed Aryl Sulfide Synthesis through an Aryl Exchange Reaction  
Isshiki et al. (2021) | *Journal of the American Chemical Society* | DOI: 10.1021/jacs.1c04215



Ni-Catalyzed Aryl Sulfide Synthesis through an Aryl Exchange Reaction. Credit: Waseda University


Aryl sulfide, an aromatic compound in which sulfur is attached to an aryl (a functional group derived from an aromatic ring), is found in biologically active materials effective against asthma, Alzheimer's disease, and cancer. As a result, chemists have shown a lot of interest in synthesizing aryl sulfides. Traditionally, carbon-sulfur (C-S) bond

formation reactions between thiols and aryl electrophiles catalyzed by transition metals have been employed for aryl sulfide synthesis because of their high reliability. However, thiols have an unpleasant smell and are toxic. Could there be a way to synthesize aryl sulfides that avoids the use of thiols?

A team of chemists from Waseda University, Japan, led by Professor Junichiro Yamaguchi addresses this question in a recent study published in the *Journal of the American Chemical Society*, and has come up with a technique that gets the job done without thiols. The team took the hint from a previous study where they used a nickel catalyst to synthesize aromatic esters from two [aromatic compounds](#). "In 2020, we developed the first ester synthesis method using [aromatic ring](#) exchange reactions and decided to apply the knowledge gained from this reaction to realize a thiol-free [sulfide](#) synthesis," explains Yamaguchi, speaking of the origin of the study.

Against this backdrop, the team set out to synthesize aryl sulfides and aromatic esters. They started out by reacting 4-tolyl sulfide and 4-phenylbenzoate in presence of a nickel catalyst and found that the desired aryl sulfide was synthesized in the presence of a ligand, dcypt, and a zinc additive,  $\text{Zn}(\text{OAc})_2$ .

Encouraged by the results, the team moved on to investigate the mechanism of the reaction. They reacted the aryl sulfide with the [nickel catalyst](#) ( $\text{Ni}(\text{cod})_2$ ) and the ligand, dcypt, and observed the formation of a nickel complex consisting of the catalyst, the ligand, and the aryl sulfide. This nickel complex could react with the aromatic ester to form a pair of nickel complexes which could react with each other to form the desired aryl sulfide.



A team of chemists from Waseda University, Japan, set out to synthesize aryl sulfides using aryl sulfides and aromatic esters after developing an ester synthesis involving an aromatic ring exchange. Credit: Junichiro Yamaguchi from Waseda University, Japan

From these reactions, the team concluded that the Ni-catalyzed aryl sulfide synthesis occurred in a sequence of steps. Initially, the aryl sulfide and aromatic esters underwent simultaneous oxidative addition reactions to the nickel/ligand catalyst forming nickel complexes. These nickel complexes could engage in an aryl exchange reaction to form a set of nickel intermediates. This was then followed by the reductive elimination of the intermediates to regenerate the Ni/ligand catalyst and form the desired aryl sulfide.

The reductive elimination of the [nickel](#) intermediates, however, reduced the yield of the aryl sulfide. To combat this, the team employed 2-pyridyl sulfide, which accelerated this limiting step, improving the

yield. Additionally, the synthesis method worked with several aryl electrophiles such as aromatic esters, arenol derivatives and aryl halides.

"The sulfide synthesis method developed can proceed for a variety of complex bioactive compounds such as probenecid, flavone, estrone, phenylalanine, umbelliferone, and  $\beta$ -isocupreidine derivatives," comments an excited Yamaguchi, contemplating the prospects of their novel [synthesis](#) technique. "Furthermore, the ability to use environmentally friendly [aromatic esters](#) and phenol derivatives as feedstock and pyridyl sulfide as a sulfide agent could make this technology suitable for both laboratory and industrial-scale applications."

**More information:** Ryota Isshiki et al, Ni-Catalyzed Aryl Sulfide Synthesis through an Aryl Exchange Reaction, *Journal of the American Chemical Society* (2021). [DOI: 10.1021/jacs.1c04215](https://doi.org/10.1021/jacs.1c04215)

Provided by Waseda University

Citation: Synthesizing aryl sulfides from non-smelling, non-toxic compounds (2021, July 21) retrieved 23 April 2024 from <https://phys.org/news/2021-07-aryl-sulfides-non-smelling-non-toxic-compounds.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.