

Mapping extreme snowmelt and its potential dangers

July 12 2021, by Mikayla MacE Kelley



The map shows the greatest amount of snow loss over a two day period across the United States within a 30-year window. The largest snow loss, indicated by green and blue, occurs in the mountains of the western United States. Units are millimeters of snow mass lost per two days. Only pixels, which equate to 2.5 square miles each, with extreme snow loss (exceeding 50 mm per two days) are included. Credit: Josh Welty

Snowmelt—the surface runoff from melting snow—is an essential water



resource for communities and ecosystems. But extreme snow melt, which occurs when snow melts too rapidly over a short amount of time, can be destructive and deadly, causing floods, landslides and dam failures.

To better understand the processes that drive such rapid melting, researchers set out to map extreme snowmelt events over the last 30 years. Their findings are published in a new paper in the *Bulletin of the American Meteorological Society*.

"When we talk about snowmelt, people want to know the basic numbers, just like the weather, but no one has ever provided anything like that before. It's like if nobody told you the maximum and <u>minimum</u> temperature or record temperature in your city," said study co-author Xubin Zeng, director of the UArizona Climate Dynamics and Hydrometeorology Center and a professor of atmospheric sciences. "We are the first to create a map that characterizes snowmelt across the U.S. Now, people can talk about the record snowmelt events over each small area of 2.5 miles by 2.5 miles."

Zeng and lead study author Josh Welty, who received his <u>doctoral degree</u> under Zeng's advising, created a map that catalogs the top 10 extreme snowmelt events in terms of frequency, magnitude, temperature and precipitation over every 2.5-mile square of the U.S. between 1988 and 2017. They also used machine learning to understand how large-scale weather patterns affect extreme snow melt.

They found that in the western half of the country, winds transport water vapor from the Pacific Ocean eastward. However, in the eastern half of the country, weather patterns transport moisture primarily south to north from the Gulf of Mexico all the way to the Great Lakes and New England.



Their maps also reveal that in most cases, extreme snowmelt is caused by unusually warm temperatures. This conclusion is fairly intuitive, but a surprising finding revealed that in certain regions, particularly in the Pacific Northwest and the northeastern U.S., extreme snowmelt events are driven by rain—which is relatively warm—falling on snow. In these cases, extreme snowmelt events become immediately dangerous.

The paper outlines one such example in detail: The Oroville Dam in Butte County, California, holds the second-largest reservoir in the state. In 2017, a series of storms dropped huge amounts of warm rain on the snowcapped Sierra Nevada Mountains, resulting in rapid snowmelt that filled the dam past its brim. Spillways, which provide controlled water runoff, failed, and over 180,000 people were evacuated.

Such events might happen more often in the future, according to Zeng and Welty's findings. The researchers found only a slight increase in the frequency of such events over the 30-year period, and they didn't see a trend in terms of the magnitude of extreme <u>snowmelt</u> events. However, 30 years isn't long enough to establish a trend, said Zeng, who is also the Agnes N. Haury Endowed Chair in Environment in the UArizona Department of Hydrology and Atmospheric Sciences. That means future research will be especially important.

"This paper serves as foundation and a reference point to see if and how things will be changing in different regions over the next 10 to 15 years," Welty said.

More information: Josh Welty et al, Characteristics and Causes of Extreme Snowmelt over the Conterminous United States, *Bulletin of the American Meteorological Society* (2021). DOI: 10.1175/BAMS-D-20-0182.1



Provided by University of Arizona

Citation: Mapping extreme snowmelt and its potential dangers (2021, July 12) retrieved 3 May 2024 from <u>https://phys.org/news/2021-07-extreme-snowmelt-potential-dangers.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.