


Martian global dust storm ended winter early in the south

July 23 2021



Images of Mars under clear conditions (left) and during the 2018 Global Dust Storm (right). Credit: NASA, ESA, STScI.

A dust storm that engulfed Mars in 2018 destroyed a vortex of cold air around the planet's south pole and brought an early spring to the hemisphere. By contrast, the storm caused only minor distortions to the polar vortex in the northern hemisphere and no dramatic seasonal changes. Dr. Paul Streeter of The Open University's Faculty of Science,

Technology, Engineering and Mathematics will present the work today (23 July) at the virtual [National Astronomy Meeting](#) (NAM 2021).

Over two weeks at the beginning of June 2018, localized dust storms combined and spread to form an impenetrable blanket of dust that hid almost the entire planet's surface. The [global dust storm](#), which coincided with Mars's equinox and lasted until mid-September, proved fatal to NASA's solar-powered Opportunity rover.

Streeter and colleagues from The Open University, NASA and the Russian Academy of Sciences examined the effects of the event on the [martian atmosphere](#) by combining data from a Mars Global Climate Model with observations from the European Space Agency/Roscosmos ExoMars Trace Gas Orbiter and NASA's Mars Reconnaissance Orbiter missions.

Dr. Streeter said: "This was a perfect opportunity to investigate how global dust storms impact the atmosphere at the martian poles, which are surrounded by powerful jets of wind in winter. Since the last global storm in 2007, several new missions and instruments have arrived in Mars orbit, so the 2018 event was the most-observed to date."

Previous research has shown that high levels of dust in the atmosphere can have significant effects on polar temperatures and winds. The vortices at the winter poles also affect temperatures and the transport of air, dust, water and chemicals, so their disruption could mean substantial changes in the martian atmosphere.

The team found that the 2018 storm had profoundly different effects in each hemisphere. At the [south pole](#), where the vortex was almost destroyed, temperatures rose and wind speeds fell dramatically. While the vortex may have already been starting to decay due to the onset of spring, the dust storm appears to have had a decisive effect in ending

winter early.

The northern [polar vortex](#), by contrast, remained stable and the onset of autumn followed its usual pattern. However, the normally elliptical northern vortex was changed by the [storm](#) to become more symmetrical. The researchers link this to the high dust content in the atmosphere suppressing atmospheric waves caused by the extreme topography in the [northern hemisphere](#), which has volcanoes over twice as tall as Mount Everest and craters as deep as terrestrial mountains.

Dr. Streeter added: "Global dust storms at equinox may enhance transport into the southern pole due to the diminished vortex, while the more robust northern vortex continues to act as an effective barrier. If this pattern for global [dust storms](#) holds over the course of the thousands of years that Mars maintains this particular axial tilt, it has implications for how [dust](#) is deposited at the north and south poles and our understanding of the planet's climate history."

Provided by Royal Astronomical Society

Citation: Martian global dust storm ended winter early in the south (2021, July 23) retrieved 11 July 2024 from <https://phys.org/news/2021-07-martian-global-storm-winter-early.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.