


Researchers reveal robust ethane-trapping porous organic cage for efficient ethylene purification application

July 9 2021, by Liu Jia



Credit: Prof. YUAN's Group

The removal of ethane (C₂H₆) from its analogous ethylene (C₂H₄) is of great importance in the petrochemical industry, and is highly challenging due to their similar physicochemical properties. The use of emerging porous organic cage (POC) materials for C₂H₆/C₂H₄ separation is still in

its infancy.

Calix[4]resorcinarene, a kind of macrocyclic cavitand with an intrinsic cavity and eight polar upper-rim phenolic groups, has been documented as an excellent building block to construct cage compounds with tunable cavities for encapsulating various guest molecules.

In a study published in *Nature Communications*, the research group led by Prof. Yuan Daqiang from Fujian Institute of Research on the Structure of Matter of the Chinese Academy of Sciences reported that a [6+12] octahedral calix[4]resorcinarene-based POC (CPOC-301) is an excellent C_2H_6 -selective material, and can be used as a robust absorbent to directly afford high-purity C_2H_4 from C_2H_6/C_2H_4 mixture.

The researchers prepared CPOC-301 via self-assembly of tetraformylresorcin[4]arene and p-phenylenediamine under mild conditions.

Single-crystal X-ray diffraction revealed that CPOC-301 has a truncated octahedron structure, with eight trigonal ports having edge lengths reaching about 12 Å, and a large cavity with inner diameter as well as volume that respectively reach 16.8 Å and 4270 Å³. The solid-state packing of CPOC-301 suggested that it possesses a one-dimensional channel, with a diameter of ~7 Å, viewed from [001] direction.

The nitrogen (N_2) gas adsorption isotherm of CPOC-301 showed a typical type I curve with a small fraction of a type IV adsorption behavior. The maximum N_2 adsorption is 670 cm³ g⁻¹, and the calculated Brunauer–Emmett–Teller (BET) surface area of CPOC-301 is up to 1962 m² g⁻¹. Moreover, CPOC-301 exhibited preferential adsorption of C_2H_6 over C_2H_4 at room temperature.

The ideal adsorbed solution theory (IAST) calculation result

demonstrated that C_2H_6/C_2H_4 selectivity range for CPOC-301 is from 1.3 to 1.4 at 293 K. The breakthrough curves further proved that CPOC-301 can efficiently realize the complete separation of C_2H_4 from C_2H_6/C_2H_4 mixtures, and its separation performance does not obviously change within seven continuous cycles.

Molecular modeling studies suggested the exceptional C_2H_6 selectivity is due to the suitable resorcin[4]arene cavities in CPOC-301, which form more multiple C–H $\cdots\pi$ hydrogen bonds with C_2H_6 than with C_2H_4 guests.

This study sheds light on the design and synthesis of POCs based on supramolecular cavitands as "porous additives" in column and membrane separation applications for industrially important gases in the future.

More information: Kongzhao Su et al, Efficient ethylene purification by a robust ethane-trapping porous organic cage, *Nature Communications* (2021). [DOI: 10.1038/s41467-021-24042-7](https://doi.org/10.1038/s41467-021-24042-7)

Provided by Chinese Academy of Sciences

Citation: Researchers reveal robust ethane-trapping porous organic cage for efficient ethylene purification application (2021, July 9) retrieved 24 April 2024 from <https://phys.org/news/2021-07-reveal-robust-ethane-trapping-porous-cage.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.