


# Symbionts sans frontieres: Bacterial partners travel the world

July 12 2021

---



Fluorescence microscopy reveals that lucinid gills are packed with symbionts. Lucinids host them in specialized cells called bacteriocytes. Bacterial symbionts are labeled in green and magenta, host nuclei in gold. Credit: Lukas Leibrecht

This pandemic year has restricted international travel for humans. Not so for some microscopic bacteria in the ocean: They partner up with clams living in the sand beneath the shimmering waters of coastal habitats throughout the globe. According to research by scientists from the Max Planck Institute for Marine Microbiology in Bremen and the University of Vienna now published in *PNAS*, the bacterial symbionts living in lucinid gills travel the world without borders.

The Lucinidae family, lucinids for short, comprises approximately 500 living species of bivalves. They are at least 400 million years old, according to [fossil records](#), and have managed to colonize a wide variety of habitats, from beautiful beaches to the abyssal depths untouched by the sun over a kilometer below the sea surface. Their ability to thrive in a wide variety of habitats is made possible by their 'partner in crime,' a sulfur-oxidizing bacterial [symbiont](#) that utilizes hydrogen sulfide, better known as 'rotten egg gas,' as an energy source to power primary production. This process is not unlike photosynthesis used by plants, yet not dependent on sunlight, and generates enough sugars to feed both the symbiont and the lucinids themselves.

## **Striking up partnerships from near or far**

Finding a suitable partner out in the wild is a matter of life and death for lucinids. They have to pick up their bacterial partners at a very early life stage when they settle in the sediment after their larval stage. From this

time on, they rely on their [bacterial symbionts](#) for nutrition. However, bacterial cells are miniscule and the oceans are awash with a multitude of possible candidates. Typically, animals that rely so heavily on bacteria are expected to strike up partnerships with local residents. These microbes are likely to work best under the unique conditions of their local habitats. A new study based on metagenomic analyses of symbiotic bacteria in lucinids now reveals that this is not always the case: Some bacterial symbionts travel the globe and are true cosmopolitans.

## Globally distributed symbionts

"Using state-of-art DNA-sequencing and genome assembling, we discovered that a single bacterial symbiont species was the most abundant symbiont in eight lucinid species spanning three oceans—the Atlantic, Pacific and Indian Oceans—across the tropics of both hemispheres," said Laetitia Wilkins from the Max Planck Institute for Marine Microbiology in Bremen, Germany, shared first author of the publication together with Jay Osvatic from the University of Vienna, Austria. "These symbionts are virtually all over the place." No other known symbiont is so successful at dispersal and establishing symbioses with lucinids, the researchers report. They named it *Candidatus Thiodiazotropha taylori*—"to acknowledge the wisdom of John Taylor from the Natural History Museum in London, who has devoted 25 years of his life to studying lucinid biology and taxonomy," Osvatic said.

"This unexpected finding challenges previous concepts that symbionts are acquired locally. It suggests that lucinid symbionts are much more mobile," Osvatic said. The remarkable flexibility in this partnership is advantageous to both host and symbiont, as it increases the likelihood of locating a compatible partner across diverse habitats all over the globe. Prior to this study, lucinid research has mainly been carried out in easily accessible locations. Now for the first time the team around Wilkins and Osvatic presents an expanded, global dataset that has led to and will

continue to facilitate [new discoveries](#) and show how distant habitats might be connected.




Lucinids are the most species rich and widely distributed family of marine bivalves hosting bacterial endosymbionts. In this picture, a large specimen of *Ctena imbricatula* is checking out its environment with its foot that it can enlarge ten times its body size. Credit: Laetitia Wilkins

### **Scientists collaborating to find collaborating organisms**

Just like the relationships between symbionts and lucinid clams, this

discovery would not have been possible without the scientists reaching out and forming collaborations across the world. "Our contacts (and now friends) across the world have given us access to an unprecedented diversity of lucinids, both direct from the beaches and from museums across the world," said Benedict Yuen from the University of Vienna, senior author of the paper. "We were given access to a wide variety of lucinid samples at the Natural History Museum in London through John Taylor. Samples were also collected personally by our team and collaborators Matthieu Leray in Panama, Yolanda Camacho in Costa Rica, Olivier Gros in Guadeloupe and Jan A. van Gils in Mauritania."



This is where the investigated clams like to live: Lucinid habitat characterized by sandy patches within seagrass (*Thalassia testudinum*) beds in Bocas del Toro,

Panama. Credit: Laetitia Wilkins

### **Also discovered: Two new species in cozy togetherness**

Moreover, the extensive data collection of Wilkins, Osvatic and their team resulted in the discovery of two new lucinid symbionts, which have now been described and named after Miriam Weber and Christian Lott, both former researchers from the Max Planck Institute for Marine Microbiology in Bremen. These symbionts—now known as *Thiodioazotropha weberae* and *lotti*—are found in the clam *Loripes orbiculatus* on the Italian island of Elba, where the symbionts peacefully co-exist in the gills of the same host. "Before genomic analyses were used, it was assumed that each clam hosts only one species of symbionts," Wilkins explained. "However, many clams on Elba harbor two symbiont species. Miriam and Christian discovered this clam population in the bay of Fetovaia and it is thanks to them that we could amass a very powerful dataset on this symbiosis."

Next, the researchers want to find out how the symbionts travel. "They leave their bivalve home to traverse the globe," added co-senior author Jillian Petersen. "Both beneficial symbionts such as *Candidatus T. taylora* but also pathogens can disperse in the environment, but we usually don't know how."

**More information:** Jay T. Osvatic et al., "Global biogeography of chemosynthetic symbionts reveals both localized and globally distributed symbiont groups," *PNAS* (2021).

[www.pnas.org/cgi/doi/10.1073/pnas.2104378118](http://www.pnas.org/cgi/doi/10.1073/pnas.2104378118)

Provided by Max Planck Society

Citation: Symbionts sans frontieres: Bacterial partners travel the world (2021, July 12) retrieved 1 May 2024 from

<https://phys.org/news/2021-07-symbionts-sans-frontieres-bacterial-partners.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.