


Very-high energy gamma-ray emission detected from blazar TXS 1515–273

July 27 2021, by Tomasz Nowakowski





Multi-wavelength light curve of TXS 1515–273. Credit: MAGIC Collaboration, 2021.

An international team of astronomers has carried out multi-wavelength observations of a blazar known as TXS 1515–273; they detected veryhigh energy (VHE) gamma-ray emission from this source for the first time. The finding is detailed in a paper published July 20 on arXiv.org.

Blazars, classified as members of a larger group of active galaxies that host active galactic nuclei (AGN), are the most numerous extragalactic gamma-ray sources. Their characteristic features are relativistic jets pointed almost exactly toward the Earth. Based on their optical emission properties, astronomers divide blazars into two classes: flat-spectrum radio quasars (FSRQs) that feature prominent and broad optical emission lines, and BL Lacertae objects (BL Lacs), which do not.

At a redshift of 0.1285, TXS 1515–273 is a <u>blazar</u> of BL Lac class. Previous observations of this object found that it has a photon index of about 2.0, which suggests that it may be an extreme high frequency-peaked blazar (EHBL). In general, EHBLs are excellent targets to search for new extragalactic VHE <u>gamma-ray emission</u>.

In the beginning of 2019 a flaring activity in the high-energy (HE) gamma-ray band started from TXS 1515–273. Hoping to detect VHE gamma-ray emission, a team led by researchers from the Major Atmospheric Gamma-ray Imaging Cherenkov (MAGIC) telescope collaboration commenced a multi-wavelength observational campaign of this source.

"A MWL [multi-wavelength] campaign on TXS 1515–273 was organized at the end of February 2019. During the flaring activity, the



source was observed in different energy bands, ranging from radio to VHE gamma rays. Simultaneous or quasi-simultaneous observations were carried out by KVA, Swift, XMM, NuSTAR and MAGIC in order to investigate the location of the SED [spectral energy distribution] peaks and look for evidence of extreme behavior during the flare," the astronomers wrote in the paper.

MAGIC observed TXS 1515–273 between February 27 and March 5, 2019, which resulted in a significant detection of this blazar in the VHE gamma-ray band. The average integral VHE flux of TXS 1515–273 was measured to be approximately 6 percent of the Crab nebula flux above 400 GeV.

The astronomers detected an hour-scale variability in the X-ray fluxes of TXS 1515–273 and a clear change in the spectral shape between XMM-Newton and NuSTAR observations. The data from the X-ray variability timescales allowed the team to determine the synchrotron peak frequency of the blazar. The results suggest that the source is a flaring high, but not extreme, synchrotron peaked object.

Studying the jet of TXS 1515–273, the researchers found no new components or moving components. This is consistent with what is generally seen in other TeV blazars with high synchrotron peak frequency.

As part of the study, the scientists also modeled the broadband SED from radio to VHE gamma rays.

"We applied a simple one-zone model, which could not reproduce the radio emission and the shape of the optical emission, and a two-component leptonic model with two interacting components, enabling us to reproduce the emission from radio to VHE band," the authors of the paper explained.



More information: First detection of VHE gamma-ray emission from TXS 1515–273, study of its X-ray variability and spectral energy distribution, arXiv:2107.09413 [astro-ph.HE] arxiv.org/abs/2107.09413

© 2021 Science X Network

Citation: Very-high energy gamma-ray emission detected from blazar TXS 1515–273 (2021, July 27) retrieved 13 March 2024 from https://phys.org/news/2021-07-very-high-energy-gamma-ray-emission-blazar.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.