


# Zero-dimensional molecular sieve membranes to enhance gas separation selectivity

July 21 2021



Schematic illustration of SAM assembly. Credit: DICP

Classical molecular sieve membranes, with 3-D microparticles and 2-D nanosheets as primary building blocks, are promising in chemical separation.

Separation within such membranes relies on molecular movement and transport through their intrinsic or artificial nanopores. Since the weak connections by nature between the neighboring "bricks" usually result in intercrystalline gaps in membranes, the prevailing selectivity for classical

molecular sieve membranes is moderate.

Recently, a research group led by Prof. Yang Weishen and Dr. Ban Yujie from the Dalian Institute of Chemical Physics (DICP) of the Chinese Academy of Sciences (CAS) proposed zero-dimensional molecular sieve membranes that could enhance the separation selectivity of hydrogen ( $H_2$ ) and carbon dioxide ( $CO_2$ ).

The study was published in *Angewandte Chemie International Edition* on July 16.

"Zero-dimensional molecules, as primary building blocks in the proposed membrane, have the potential to absolutely eliminate intercrystalline gaps in membranes," said Dr. Ban.

The researchers fabricated the zero-dimensional molecular sieve [membrane](#) by orderly assembling zero-dimensional 2-methylimidazole (mim) molecules into unprecedented supramolecule array membranes (SAMs) through solvent-free vapor processing on a metal-organic framework.

In SAMs, the "zero-dimensional building blocks" together with supramolecule interactions resulted in the absence of the intercrystalline gaps, which guaranteed an effective mass-transfer through intermolecular spacings instead of an undesirable leakage through non-selective gaps.

In contrast to the classical transport through nanopores of membranes, selective transport through the intermolecular spacing of mim (~0.30 nm) was realized within SAMs, yielding an extremely precise sieving of  $H_2$  from  $CO_2$ . The  $H_2/CO_2$  selectivity was one order of magnitude higher than selectivities of the state-of-the-art classical membranes.

"Our study opens the door to create a variety of SAMs to distinguish the subtle size/shape differences of a pair of gas molecules," said Prof. Yang. "In the future, we will tailor the intermolecular spacing, control the assembly process, and enable a wide range of application of SAMs to energy-efficient [chemical separation](#) processes."

**More information:** Weishen Yang et al, High-selective Supramolecule Array Membrane Made of Zero-dimensional Molecules for Gas Separation, *Angewandte Chemie International Edition* (2021). [DOI: 10.1002/anie.202108185](https://doi.org/10.1002/anie.202108185)

Provided by Chinese Academy of Sciences

Citation: Zero-dimensional molecular sieve membranes to enhance gas separation selectivity (2021, July 21) retrieved 3 May 2024 from <https://phys.org/news/2021-07-zero-dimensional-molecular-sieve-membranes-gas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.